


Accepted Manuscript

Alcohol consumption and prevalence of erectile dysfunction in Japanese patients with type 2 diabetes mellitus: baseline data from The Dogo Study

Shinya Furukawa, Takenori Sakai, Tetsuji Niiya, Hiroaki Miyaoka, Teruki Miyake, Shin Yamamoto, Kotatsu Maruyama, Teruhisa Ueda, Keiko Tanaka, Hidenori Senba, Yasuhiko Todo, Masamoto Torisu, Hisaka Minami, Morikazu Onji, Takeshi Tanigawa, Bunzo Matsuura, Yoichi Hiasa, Yoshihiro Miyake



PII: S0741-8329(15)30020-3

DOI: [10.1016/j.alcohol.2016.07.006](https://doi.org/10.1016/j.alcohol.2016.07.006)

Reference: ALC 6613

To appear in: *Alcohol*

Received Date: 14 April 2015

Revised Date: 18 March 2016

Accepted Date: 21 July 2016

Please cite this article as: Furukawa S., Sakai T., Niiya T., Miyaoka H., Miyake T., Yamamoto S., Maruyama K., Ueda T., Tanaka K., Senba H., Todo Y., Torisu M., Minami H., Onji M., Tanigawa T., Matsuura B., Hiasa Y. & Miyake Y., Alcohol consumption and prevalence of erectile dysfunction in Japanese patients with type 2 diabetes mellitus: baseline data from The Dogo Study, *Alcohol* (2016), doi: [10.1016/j.alcohol.2016.07.006](https://doi.org/10.1016/j.alcohol.2016.07.006).

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Alcohol consumption and prevalence of erectile dysfunction in
Japanese patients with type 2 diabetes mellitus: baseline data
from The Dogo Study**

Shinya Furukawa^a, Takenori Sakai^b, Tetsuji Niiya^c, Hiroaki Miyaoka^d, Teruki Miyake^e, Shin Yamamoto^e, Kotatsu Maruyama^f, Teruhisa Ueda^g, Keiko Tanaka^a, Hidenori Senba^c, Yasuhiko Todo^e, Masamoto Torisu^h, Hisaka Minamiⁱ, Morikazu Onji^j, Takeshi Tanigawa^f, Bunzo Matsuura^k, Yoichi Hiasa^e, Yoshihiro Miyake^a

^aDepartment of Epidemiology and Preventive Medicine, Ehime University Graduate School of Medicine, Ehime, Japan

^bDepartment of Internal Medicine, Yawahama General City Hospital, Ehime, Japan

^cDepartment of Internal Medicine, Matsuyama Shimin Hospital, Ehime, Japan

^dDepartment of Internal Medicine, Saiseikai Matsuyama Hospital, Ehime, Japan

^eDepartment of Gastroenterology and Metabology, Ehime University Graduate School of Medicine, Ehime, Japan

^fDepartment of Public Health, Juntendo University School of Medicine, Tokyo, Japan

^gDepartment of Internal Medicine, Ehime Central Hospital, Ehime, Japan

^hDepartment of Internal Medicine, Saiseikai Saijo Hospital, Ehime, Japan

ⁱDepartment of Internal Medicine, Ehime Niihama Hospital, Ehime, Japan

^jDepartment of Internal Medicine, Saiseikai Imabari Hospital, Ehime, Japan

^kDepartment of Lifestyle-related Medicine and Endocrinology, Ehime University Graduate School of Medicine, Ehime, Japan

Corresponding author:

Shinya Furukawa, MD, Ph.D.

Department of Public Health, Ehime University Graduate School of Medicine
Shitsukawa, Toon, Ehime 791-0295, Japan

Telephone: +81 89 960 5283

Fax: +81 89 690 5384

Download English Version:

<https://daneshyari.com/en/article/7501592>

Download Persian Version:

<https://daneshyari.com/article/7501592>

[Daneshyari.com](https://daneshyari.com)