Accepted Manuscript

Title: Using Simulated Family Presence to Decrease Agitation in Older Hospitalized Delirious Patients: A Randomized Controlled Trial

Authors: Christine M. Waszynski, Kerry A. Milner, Ilene Staff, Sheila L. Molony

PII: DOI: Reference: S0020-7489(17)30225-0 https://doi.org/10.1016/j.ijnurstu.2017.09.018 NS 3028

To appear in:

 Received date:
 5-3-2017

 Revised date:
 26-8-2017

 Accepted date:
 29-9-2017

Please cite this article as: Waszynski, Christine M., Milner, Kerry A., Staff, Ilene, Molony, Sheila L., Using Simulated Family Presence to Decrease Agitation in Older Hospitalized Delirious Patients: A Randomized Controlled Trial.International Journal of Nursing Studies https://doi.org/10.1016/j.ijnurstu.2017.09.018

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



ACCEPTED MANUSCRIPT

Title: Using Simulated Family Presence to Decrease Agitation in Older Hospitalized Delirious

Patients:

A Randomized Trial

Authors: Christine M. Waszynski DNP, RN, GNP-BC¹ Corresponding author

Kerry A. Milner, DNSc, RN²

Ilene Staff, PhD³

Sheila L.Molony, PhD, RN, GNP-BC⁴

- Department of Geriatric Medicine and Department of Nursing, Hartford Hospital 80 Seymour Street Hartford, CT USA 06012 Cwaszynski163@gmail.com
- College of Nursing, Sacred Heart University 5151 Park Ave Fairfield, CT USA 06825 Milnerk@sacredheart.edu
- Department of Research, Hartford Hospital 80 Seymour Street Hartford, CT USA 06102 Ilene.staff@hhchealth.org
- School of Nursing, Quinnipiac University 275 Mt Carmel Ave Hamden, CT USA 06518 Shelia.molony@gmail.com

Abstract

Background: Simulated family presence has been shown to be an effective nonpharmacological

intervention to reduce agitation in persons with dementia in nursing homes. Hyperactive or

mixed delirium is a common and serious complication experienced by hospitalized patients, a

Download English Version:

https://daneshyari.com/en/article/7515144

Download Persian Version:

https://daneshyari.com/article/7515144

Daneshyari.com