



## CASE REPORT

# Isolated double chambered right ventricle as a rare cause of sudden death in infancy

W. Thaljawi\*, M. Belhadj, Y. Chkirben, S. Bouslema, M. Jedidi, T. Masmoudi, M.K Souguir, M.T. Yaâcoubi, M. Ben Dhiab

Department of Legal Medicine, HUC Farhat HACHED, 4000 Sousse, Tunisia

Received 23 August 2015; revised 4 April 2016; accepted 27 April 2016

## KEYWORDS

Sudden death;  
Congenital heart disease;  
Double chambered right  
ventricle;  
Autopsy

**Abstract** Isolated double-chambered right ventricle (DCRV) is a rare form of congenital heart disorder in which the right ventricle is divided into a high-pressure inlet portion and a low-pressure outlet portion by an anomalous muscle bundle. Rarely, autopsy cases describing this congenital heart disease have been reported in the world literature.

We report the case of a 50-day-old male who died seven days after been suffering from fever and dyspnea. The autopsy revealed essentially an isolated double-chambered right ventricle (DCRV) and an important pulmonary edema.

In this paper, we present the first reported case of sudden unanticipated death in infancy due to isolated double-chambered right ventricle.

© 2016 The International Association of Law and Forensic Sciences (IALFS). Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

One of the leading causes of prenatal and infant death from congenital malformations is congenital heart disease (CHD).<sup>1</sup> Its incidence in different studies varies from 4/1000 to 50/1000 live births.<sup>2</sup> A recent report from the United Kingdom Northern Congenital Abnormality Survey showed that 10% of deaths in this pediatric cohort with at least one congenital anomaly were associated with CHD.<sup>3</sup>

Double-chambered right ventricle is a rare congenital heart disease which consists of a cardiac disease of the right ventricular outflow tract obstruction characterized by an anomalous muscle bundles (AMB) that divides the right ventricle into

two chambers: a high-pressure inflow chamber and a low-pressure outflow chamber.

We report an unusual cause of death in a 50-day-old male infant showing a double-chambered right ventricle in the autopsy.

## 2. Case report

A male neonate born at 35 weeks after a well attended pregnancy with a normal vaginal delivery to a 25 year old primigravida mother suffering from toxemia of pregnancy. Apgar scores were 8 and 9 at 5 and 10 min, respectively. Birth weight was 2800 g. The initial clinical exam was normal.

On day 43 of life he was suffering from fever and dyspnea. A chest film was required showing no abnormal features. CRP was little high and hemogram test and electrolytes were normal.

\* Corresponding author.

Peer review under responsibility of The International Association of Law and Forensic Sciences (IALFS).

<http://dx.doi.org/10.1016/j.ejfs.2016.04.005>

2090-536X © 2016 The International Association of Law and Forensic Sciences (IALFS). Production and hosting by Elsevier B.V.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

In view of sepsis, antibiotics were started. The condition continued to be critical and the clinical status evolved to the worst.

Although he was transferred to an intensive care unit, the baby deceased on his 50th day of life.

Because there was a dubiety regarding the manner of death, a forensic autopsy was requested by the district attorney almost 17 h after death.

The deceased was a well-developed, well-nourished, 3.664 g, 54 cm male infant without any evidence of trauma.

The external examination revealed a lip and subungual cyanosis, some bloody secretions from nostrils, further there was no visible injury or congenital malformations in the external examination of the body.

On autopsy, the 58 g heart (the expected weight is  $22 \text{ g} \pm 10 \text{ g}$ ) was enlarged and hypertrophied and had smooth epicardial surfaces with a few petechiae. It measured  $3.5 \text{ cm} \times 5 \text{ cm} \times 7.5 \text{ cm}$ .

The great vessels of the heart had no anomalies or developmental defects. The coronary system was left-dominant, with a normal caliber and without any atherosclerotic lesions. On the opening of the heart, the left ventricular wall was 0.83 cm in thickness and of mottled appearance and the right ventricle was widely hypertrophic; 0.43 cm in thickness (Table 1). We notified the presence of a muscle bundle measured 1.1 cm in large that protrudes from the right ventricular free wall to the interventricular septum (Fig. 1). It divides the right ventricle into two chambers.

No apparent sclerosis of the cardiac valves and no stenoses, thrombi, or anomalies of the coronary arteries were noted. The foramen ovale was closed, and we could not find any other cardiac anomalies.


The 70 g left lung and 85 g right lung had smooth pleural surfaces with multiple scattered petechiae and congested parenchyma.

Bronchi contained thick bronchial secretions and pulmonary arteries were free. The pressure of the pulmonary sections showed an issue of purulent secretions associated with massive edema.

The rest of organs, such as the brain, the liver, and the kidneys, were extremely congested without any other macroscopic abnormality.

A histological examination was performed with hematoxylin and eosin stain.

The endocardium showed pronounced thickening with laminar deposition of elastic and collagen fibers prevailing at the right ventricle. There was no fibrosis or necrosis. The conduction system had no significant lesions. The section of the right ventricular myocardium demonstrated marked myocyte hypertrophy and disorganization. The muscle bundle



**Figure 1** (a) and (b) The right ventricle (RV) is divided into two chambers by a pseudo septum.

dividing the right ventricle was composed by whorled myocardial fascicles. The myocardial cells had hypertrophic nuclei (Fig. 2).

Evidence of respiratory infection and severe congestion were disclosed by histology. The different fragments of lung have, roughly, a preserved architecture. It shows a suppurative exudate rich in neutrophils (Fig. 3a). The alveoli's lumen contains focally edematous lesion and is occupied by a dense cell population, consisting mainly of neutrophils and some macrophages ("leukocytic alveolitis") (Fig. 3b).

Other organs had no significant pathologic findings except for non-specific congestion.

**Table 1** Data concerning heart weight and ventricular wall thicknesses in our case comparing with lower-upper 95% confidence limits (tables from Scholz et al, 1988.<sup>19</sup>)

Organes	Our case	Lower-upper 95% confidence limits
Heart weight (g)	58 g	[12 g–33 g]
Left ventricular wall thicknesses (cm)	0.82 cm	[0.21 cm–0.84 cm]
Right ventricular wall thicknesses (cm)	0.43 cm	[0.06 cm–0.26 cm]
Ventricular wall thicknesses (cm)	0.92 cm	[0.22 cm–0.93 cm]

Download English Version:

<https://daneshyari.com/en/article/7531221>

Download Persian Version:

<https://daneshyari.com/article/7531221>

[Daneshyari.com](https://daneshyari.com)