



ORIGINAL ARTICLE

Aortic dilatation after tetralogy of Fallot repair: A ghost from the past or a problem in the future?

Cristina Cruz ^{a,*}, Teresa Pinho ^a, Vânia Ribeiro ^a, Cláudia Camila Dias ^b,
José Silva Cardoso ^a, Maria Júlia Maciel ^a

^a Department of Cardiology, Centro Hospitalar São João, Faculty of Medicine, University of Porto, Porto, Portugal

^b Department of Community Medicine, Information and Decision in Health, Faculty of Medicine, University of Porto, Porto, Portugal

Received 17 April 2017; accepted 1 October 2017

KEYWORDS

Aortic dilatation;
Aortic elasticity;
Tetralogy of Fallot;
Transthoracic
echocardiography

Abstract

Introduction and Aims: Intrinsic aortopathy can lead to dilatation late after tetralogy of Fallot (TOF) repair. Its extent and prevalence are not known. We aimed to assess aortic dimensions and elasticity and to find predictors of aortic dilatation.

Methods: A total of 126 adults were prospectively included after TOF repair and compared to 63 gender- and age-matched controls. Transthoracic echocardiography was used to assess aortic diameters at the level of the sinuses of Valsalva and ascending aorta and aortic dilatation was defined as z-score >+2. M-mode parameters of the ascending aorta were used to calculate strain, distensibility and stiffness index.

Results: TOF patients (mean age 30±9 years; 52% male) had a complete repair at a median age of five (2-49) years; mean follow-up time since repair was 23±7 years. The prevalence of aortic dilatation at the sinuses of Valsalva and ascending aorta was 29% and 24%, respectively. Compared to controls, TOF patients had a higher ascending aorta z-score, lower strain (6.4% [0.0-61.5] vs. 15.2% [0.0-45.0], p<0.01) and higher stiffness index (7.3 [0.8-23.6] vs. 3.1 [0.9-14.1], p<0.01). On multivariate analysis male gender was strongly associated with sinuses of Valsalva dilatation (odds ratio 6.3, 95% confidence interval 1.5-26.3, p=0.01).

Abbreviations: 2D, two-dimensional; AAo, ascending aorta; AAoZ, ascending aorta z-score; Ad, aortic diastolic diameter; Ao, aortic; AoZ, Ao root z-score; As, aortic systolic diameter; BP, blood pressure; BSA, body surface area; CI, confidence interval; D, distensibility; DBP, diastolic blood pressure; GUCH, grown-up congenital heart; LV, left ventricular; LVEF, left ventricular ejection fraction; OR, odds ratio; PP, pulse pressure; SAC, systemic arterial compliance; SBP, systolic blood pressure; SD, standard deviation; SoV, sinuses of Valsalva; STJ, sinotubular junction; SV, stroke volume; SVI, stroke volume index; TOF, tetralogy of Fallot; TTE, transthoracic echocardiography.

* Corresponding author.

E-mail address: mchristina.cruz@hsjoao.min-saude.pt (C. Cruz).

<https://doi.org/10.1016/j.repc.2017.10.014>

0870-2551/© 2018 Sociedade Portuguesa de Cardiologia. Published by Elsevier España, S.L.U. All rights reserved.

Conclusions: The prevalence of aortic dilatation late after TOF repair is significant, with a larger and stiffer ascending aorta. Male gender appears to influence aortic root dilatation. This aortopathy requires careful follow-up in order to prevent future complications.
© 2018 Sociedade Portuguesa de Cardiologia. Published by Elsevier España, S.L.U. All rights reserved.

PALAVRAS-CHAVE

Dilatação aórtica;
Elasticidade aórtica;
Tetralogia de Fallot;
Ecocardiografia
transtorácica

Dilatação da aorta na tetralogia de Fallot operada: um fantasma do passado ou um problema no futuro?

Resumo

Introdução e objetivos: Uma possível aortopatia intrínseca poderá condicionar dilatação tardia da aorta na tetralogia de Fallot operada. A sua prevalência e extensão não estão definidas. Pretendemos avaliar as dimensões e a elasticidade da aorta e encontrar preditores da dilatação aórtica.

Métodos: Incluímos prospectivamente 126 adultos operados a tetralogia de Fallot e comparamos com 63 controlos. Avaliamos por ecocardiografia transtorácica os diâmetros dos seios de Valsalva e da aorta ascendente e definimos dilatação aórtica para z -score $> +2$. Parâmetros modo M da aorta ascendente foram usados para calcular strain, distensibilidade e índice de rigidez.

Resultados: Doentes com tetralogia de Fallot (idade média 30 ± 9 anos; 52% homens) foram operados com uma idade mediana de 5 (2-49) anos; tempo médio de seguimento desde a cirurgia 23 ± 7 anos. A prevalência de dilatação dos seios de Valsalva e da aorta ascendente foi 29% e 24%, respetivamente. Comparado aos controlos, os doentes com tetralogia de Fallot apresentaram maior z -score da aorta ascendente, menor strain (6,4 [0,0-61,5] versus 15,2 [0,0-45,0]; $p < 0,01$) e maior índice de rigidez (7,3 [0,8-23,6] versus 3,1 [0,9-14,1]; $p < 0,01$). Na análise multivariada o sexo masculino associou-se significativamente à dilatação dos seios de Valsalva (*odds ratio* 6,3, intervalo de confiança de 95% 1,5-26,3; $p = 0,01$).

Conclusões: Na tetralogia de Fallot operada há uma prevalência significativa de dilatação tardia da aorta, com a aorta ascendente maior e mais rígida. O sexo masculino parece influenciar a dilatação da raiz da aorta. Esta aortopatia requer um seguimento cuidadoso para evitar complicações futuras.

© 2018 Sociedade Portuguesa de Cardiologia. Publicado por Elsevier España, S.L.U. Todos os direitos reservados.

Introduction

Tetralogy of Fallot (TOF) is the most common cyanotic congenital heart disease in which survival into adulthood is common. The guidelines recommend regular long-term follow-up in specialized grown-up congenital heart (GUCH) centers.¹

Major residual lesions after TOF repair occur at the level of the right ventricular outflow tract, including obstruction or aneurysm, and at the pulmonary valve due to regurgitation or stenosis. In addition, as first described by Capelli et al.,² there is an increasing awareness that aortic (Ao) dilatation can develop late after TOF repair. Interestingly, Ao dilatation is a fetal feature of TOF that tends to disappear after early surgical repair³ without a palliative systemic-to-pulmonary shunt, thus preventing long-standing volume overload on the overriding aorta. Ao dilatation may be a disorder of the past, not reflecting the current surgical era.⁴ Nevertheless, for many reasons, including anatomical and technical surgical issues, not all patients will benefit from early repair. Additionally, aortic histological abnormalities

present since infancy in TOF, especially in the Ao root and ascending aorta (AAo) vascular wall, can contribute to Ao dilatation.^{5,6} This intrinsic aortopathy can lead to Ao dilatation late after TOF repair, but its extent and prevalence are not known. We aimed to assess proximal thoracic aorta dimensions and elasticity by transthoracic echocardiography, in TOF patients and normal controls, and to find possible predictors of Ao dilatation.

Methods

Study population

Inclusion criteria

We prospectively included 126 adults (age ≥ 18 years) with repaired TOF, without pulmonary atresia or absent pulmonary valve, followed at our GUCH center between March 2011 and December 2013, and 63 gender- and age-matched healthy controls.

Download English Version:

<https://daneshyari.com/en/article/7535756>

Download Persian Version:

<https://daneshyari.com/article/7535756>

[Daneshyari.com](https://daneshyari.com)