


Contents lists available at [ScienceDirect](https://www.sciencedirect.com)

## Statistics and Probability Letters

journal homepage: [www.elsevier.com/locate/stapro](http://www.elsevier.com/locate/stapro)

# Statistical issues in radiosonde observation of atmospheric temperature and humidity profiles

A. Fassò <sup>a,\*</sup>, F. Finazzi <sup>a</sup>, F. Madonna <sup>b</sup><sup>a</sup> University of Bergamo, Department MIPE, Viale Marconi 5, Dalmine, BG, Italy<sup>b</sup> CNR-IMAA, C.da S. Loja, Tito Scalco, PZ, Italy

## ARTICLE INFO

*Article history:*

Available online xxxx

*Keywords:*Functional data  
Spatio-temporal models  
GRUAN  
RAOB

## ABSTRACT

Measurement uncertainty of atmospheric profiles obtained by radiosoundings is crucial in climate change studies. This paper shows how the understanding of geographic gaps of radiosonde networks calls for a functional approach able to handle spatio-temporal profile data, and related complexity issues are addressed.

© 2018 Elsevier B.V. All rights reserved.

## 1. Introduction

Measurement system uncertainty in climate data records (CDRs) and its impact in climate change assessment have been raised by various climate scientists and metrologists. This is especially true for observations of temperature and humidity provided by radiosoundings. Addressing this issue, the GRUAN reference measurement network (GCOS Reference Upper-Air Network, [www.gruan.org](http://www.gruan.org)) has been established, and started to provide valuable contribution to the understanding of measurement uncertainty, see [Bodeker et al. \(2016\)](#). Although GRUAN gives fully traceable measurements, its geographic and historical coverage is quite limited. For this reason climatological studies are largely based on baseline measurement networks, which have an intermediate metrological quality but a larger spatio-temporal coverage.

In perspective, integrated data sets will include both ground based and satellite observations of land, sea and atmosphere. Moreover ensembles, obtained by various simulation techniques, are of increasing importance to describe uncertainty. Hence, considering the data growth rate we are facing, the size of such data sets is fast increasing from the order of terabytes to petabytes. This frame requires a new cooperation effort to elaborate new multi-disciplinary approaches and services for different types of users. Hence, the integration of atmospheric, metrology, statistical and computer sciences is considered a fruitful route to fully exploit the available historical CDRs of several Essential Climate Variables (ECVs) collected by satellite observation platforms and by ground based networks operating at the global scale.

In this frame the uncertainty of baseline networks is a challenging issue and the present paper aims at being a step to face this problem. In particular geographic gaps of temperature and humidity radiosonde networks are discussed and the need for advanced statistical methods is illustrated.

The remainder of the paper is organized as follows. In Section 2 ongoing projects involving production and analysis of climate data sets in general and in particular radiosoundings are discussed and showed to call for new statistical developments. Section 3 considers spatio-temporal modeling for functional data in connection to network gap identification. Moreover Section 4 deepens some computational issues related to data and model size.

\* Corresponding author.

E-mail address: [alessandro.fasso@unibg.it](mailto:alessandro.fasso@unibg.it) (A. Fassò).

## 2. Data sets and projects

Copernicus is the European union's Earth observation programme ([www.copernicus.eu](http://www.copernicus.eu)). Its aim is to help in understanding how our planet and its climate are changing, the role played by human activities in these changes and how these will influence our daily lives. To do this Copernicus is involved in a complex set of systems, which collect data from multiple sources: earth observation satellites and in situ sensors such as ground stations, airborne and sea-borne sensors. It processes these data and provides users with reliable and up-to-date information through a set of services related to environmental and security issues.

In particular, the Copernicus Climate Change Service (C3S), which is operated by the European Centre for Medium-range Weather Forecasts (ECMWF, [www.ecmwf.int](http://www.ecmwf.int)), will provide comprehensive climate information covering a wide range of components of the Earth-system and timescales spanning decades to centuries. It will maximize the use of past, current and future earth observations (from in-situ and satellite observing systems) in conjunction with modeling, supercomputing and networking capabilities. This will produce a consistent, comprehensive and credible description of the past, current and future climate.

Various other projects developed in the frame of the Horizon 2020 program have been the scientific precursors of C3S. In particular Fiduceo (Fidelity and Uncertainty in climate data records from Earth Observations, [www.fiduceo.eu](http://www.fiduceo.eu)) aims at bringing insights from metrology to the observation of Earth's climate from space (Merchant et al., 2017). Moreover, GAIA-CLIM (Gap Analysis for Integrated Atmospheric ECV CLimate Monitoring, [www.gaia-clim.eu](http://www.gaia-clim.eu)) aims at understanding gaps in integrated monitoring of upper troposphere and to improve our ability to use ground-based and sub-orbital observations to characterize satellite observations for a number of atmospheric ECVs, see Thorne et al. (2017). In fact despite that satellite Earth observation technology has undoubtedly facilitated the development of global climate change research, ground-based networks such as radiosonde networks are required to identify biases and issues in the satellite CDRs. Therefore, radiosonde observations remain an essential component of the observing system of systems. In this frame, geographic gaps are characterized by poor spatial coverage of a monitoring network.

On their turn, to represent a reliable and effective reference information all of these conventional anchor data sources must be harmonized and homogenized to achieve physical consistency of the decadal time series. Homogenization is essentially change detection and adjustment of data for any kind of known and quantifiable inhomogeneities (bias, change of sensors, calibration drift, local environment changes, etc.). Homogenization methods that have been developed for radiosonde has a long history, see for example Haimberger et al. (2012), Thorne et al. (2011) and Sherwood et al. (2008). Although the statistical interpretation of these methods is very interesting, it is omitted here for brevity. Harmonization is involved with the traceable characterization of the total uncertainty budget. For example, the harmonization of temperature and humidity CDRs is one of the funded activity by C3S under contract C3S\_311a\_Lot3 (Madonna et al., 2017), and may benefit from the geographic gap analysis discussed in this paper.

## 3. Statistical issues and modeling

One of the objectives of GAIA-CLIM project is to understand the information content of ground based monitoring networks and to identify geographic gaps of these networks. We focus here on the network of the Universal RAWinsonde OBServation program or simply RAOB ([www.raob.com](http://www.raob.com)), which has a global coverage with about 2400 stations and some decades with bi-daily prevailing temporal frequency. This type of baseline networks are also involved in the C3S harmonization problem.

From the statistical point of view, Fassò et al. (2014) and Ignaccolo et al. (2015), showed that atmospheric soundings may be conveniently described as functional data using appropriate basis function expansion, at least when these data can be handled as independent replications of the same model. In case of global networks spatial and temporal correlation must be considered. Various functional models have been considered for spatio-temporal data where usually time dynamics is embedded in the functional object, see e.g. Menafoglio et al. (2013) and Mateu and Romano (2017) and references therein for recent advances in the field.

The idea of spatially correlated functional data (e.g. Delicado et al., 2010, and Ruiz-Medina, 2012) may be extended, to handle a manifold domain such as the sphere and the temporal dimension (e.g. Porcu et al., 2016). Although the idea of modeling these data as spatio-temporally correlated functional data is quite natural, from the point of view of probability theory, this object may be considered as a stochastic process defined on  $sphere \times time$  with values in a functional space. Alternatively, it may be considered as a stochastic process defined on a  $spherical\ shell \times time$  with scalar values. In both cases, the full characterization of the underlying stochastic process is still under study, including the definition of flexible families of valid covariance functions (Porcu, 2017, private communication).

Taking into account computational burden and data dimensionality, the statistical model needs to be simple enough. Various solutions for modeling large spatial data sets have been proposed including nearest neighbor models (Vecchia, 1988, and Datta et al., 2016). A first step in computation reduction is to use separable models with discrete time, possibly after adjusting for relevant trends. Following this approach, maximum likelihood estimation can be based on the EM algorithm extending the multivariate dynamic coregionalization model (Finazzi and Fassò, 2014 and Calculli et al., 2015). This can be easily done by applying a multivariate spatio-temporal model to the coefficients of the basis function expansion. In some sense this approach is close to kriging for function-valued data discussed by Delicado et al. (2010). A relevant difference is related to the smoothing factor used to obtain the basis function coefficients: while in Delicado et al. (2010) a crossvalidation

Download English Version:

<https://daneshyari.com/en/article/7548563>

Download Persian Version:

<https://daneshyari.com/article/7548563>

[Daneshyari.com](https://daneshyari.com)