

Accepted Manuscript

Application of ionic liquids for elution of bioactive flavonoid glycosides from lime fruit by miniaturized matrix solid-phase dispersion

Jing-Jing Xu, Rui Yang, Li-Hong Ye, Jun Cao, Wan Cao, Shuai-Shuai Hu, Li-Qing Peng

PII: S0308-8146(16)30158-3

DOI: <http://dx.doi.org/10.1016/j.foodchem.2016.02.012>

Reference: FOCH 18732

To appear in: *Food Chemistry*

Received Date: 22 June 2015

Revised Date: 23 December 2015

Accepted Date: 1 February 2016



Please cite this article as: Xu, J-J., Yang, R., Ye, L-H., Cao, J., Cao, W., Hu, S-S., Peng, L-Q., Application of ionic liquids for elution of bioactive flavonoid glycosides from lime fruit by miniaturized matrix solid-phase dispersion, *Food Chemistry* (2016), doi: <http://dx.doi.org/10.1016/j.foodchem.2016.02.012>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Application of ionic liquids for elution of bioactive flavonoid
glycosides from lime fruit by miniaturized matrix solid-phase
dispersion**

An abbreviated running title: Application of ILs for elution in miniaturized MSPD

Jing-Jing Xu^a, Rui Yang^b, Li-Hong Ye^c, Jun Cao^{a*}, Wan Cao^a, Shuai-Shuai Hu^a, Li-Qing Peng^a

^aCollege of Material Chemistry and Chemical Engineering, Hangzhou Normal University, Hangzhou 310036, China

^bDepartment of Surgery, University of Michigan Medical Center, Ann Arbor, Michigan 48109, United States

^cHangzhou Red Cross Hospital, Department of Traditional Chinese Medicine, Hangzhou 310003, China

* Corresponding author: Jun Cao (Dr.)

E-mail: caojun91@163.com

Tel.: +86 571 2886 7909

Fax: +86 571 2886 7909

ABSTRACT

In this work, two flavonoid glycosides (neohesperidin and naringin) in lime fruit were effectively extracted by miniaturized matrix solid phase dispersion (MSPD), followed

Download English Version:

<https://daneshyari.com/en/article/7589294>

Download Persian Version:

<https://daneshyari.com/article/7589294>

[Daneshyari.com](https://daneshyari.com)