


Accepted Manuscript

Title: Molecular characterization of biochar from five Brazilian agricultural residues obtained at different charring temperatures

Authors: Judith Schellekens, Carlos Alberto Silva, Peter Buurman, Tatiana F. Rittl, Rimena R. Domingues, Marina Justi, Pablo Vidal-Torrado, Paulo Fernando Trugilho



PII: S0165-2370(17)30933-6
DOI: <https://doi.org/10.1016/j.jaap.2018.01.020>
Reference: JAAP 4243

To appear in: *J. Anal. Appl. Pyrolysis*

Received date: 19-10-2017
Revised date: 17-1-2018
Accepted date: 18-1-2018

Please cite this article as: Judith Schellekens, Carlos Alberto Silva, Peter Buurman, Tatiana F.Rittl, Rimena R.Domingues, Marina Justi, Pablo Vidal-Torrado, Paulo Fernando Trugilho, Molecular characterization of biochar from five Brazilian agricultural residues obtained at different charring temperatures, Journal of Analytical and Applied Pyrolysis <https://doi.org/10.1016/j.jaap.2018.01.020>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Molecular characterization of biochar from five Brazilian agricultural residues obtained at different charring temperatures

Judith Schellekens^{a*}, Carlos Alberto Silva^b, Peter Buurman^c, Tatiana F. Rittl^d, Rimena R. Domingues^b, Marina Justi^b, Pablo Vidal-Torrado^a, Paulo Fernando Trugilho^e

^a *Departamento de Ciência do Solo, Escola Superior de Agricultura “Luiz de Queiroz”-ESALQ/USP, Av. Pádua Dias 11, Piracicaba, São Paulo, Brazil.*

^b *Departamento de Ciência do Solo, Universidade Federal de Lavras - UFLA, Av. Dr. Sylvio Menicucci, 1001, Lavras, Minas Gerais, Brazil.*

^c *Earth System Science Group, Wageningen University, P.O. Box 47, 6700 AA Wageningen, The Netherlands.*

^d *Institute of Meteorology and Climate Research, Atmospheric Environmental Research (IMK-IFU), Campus Alpin, Garmisch-Partenkirchen, Germany.*

^e *Departamento de Ciências Florestais, Universidade Federal de Lavras – UFLA, Av. Dr. Sylvio Menicucci, 1001, Lavras, Minas Gerais, Brazil.*

*Corresponding author. Tel.: +31 653 83 20 87.

E mail address: schellekens.j@hetnet.nl.

Highlights

- Molecular composition of biochar from five feedstocks were determined with Py-GC/MS.
- 858 pyrolysis products identified from biochars obtained at 0, 350, 450 and 750 °C (T_{char}).

Download English Version:

<https://daneshyari.com/en/article/7606386>

Download Persian Version:

<https://daneshyari.com/article/7606386>

[Daneshyari.com](https://daneshyari.com)