

Accepted Manuscript

Title: Enrichment of liposomal nanomedicines using monolithic solid phase extraction discs following preactivation with bivalent metal ion solutions

Authors: Ayako Watanabe, Koji Karasawa, Shuhei Murayama, Yoshihiro Sano, Mio Takagi, Eiichi Yamamoto, Shigeo Murata, Masaru Kato



PII: S0021-9673(18)30755-6
DOI: <https://doi.org/10.1016/j.chroma.2018.06.011>
Reference: CHROMA 359455

To appear in: *Journal of Chromatography A*

Received date: 9-2-2018
Revised date: 3-6-2018
Accepted date: 5-6-2018

Please cite this article as: Watanabe A, Karasawa K, Murayama S, Sano Y, Takagi M, Yamamoto E, Murata S, Kato M, Enrichment of liposomal nanomedicines using monolithic solid phase extraction discs following preactivation with bivalent metal ion solutions, *Journal of Chromatography A* (2018), <https://doi.org/10.1016/j.chroma.2018.06.011>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Enrichment of liposomal nanomedicines using monolithic solid phase extraction discs following preactivation with bivalent metal ion solutions

*Ayako Watanabe*¹, *Koji Karasawa*², *Shuhei Murayama*², *Yoshihiro Sano*², *Mio Takagi*¹, *Eiichi Yamamoto*³, *Shigeo Murata*^{1,4}, *Masaru Kato*^{1,2,*}

¹One-stop sharing facility center for future drug discoveries, Graduate School of Pharmaceutical Sciences, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

²Department of Bioanalytical Chemistry, School of Pharmacy, Showa University, 1-5-8 Hatanodai, Shinagawa-ku, Tokyo 142-8555, Japan

³Analytical Research, Pharmaceutical Science & Technology Unit, Medicine Development Center, Eisai Co., Ltd.

5-1-3 Tokodai, Tsukuba, Ibaraki 300-2635, Japan

⁴Laboratory of Protein Metabolism, Graduate School of Pharmaceutical Sciences, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

*Corresponding Author

Masaru Kato, Tel:+81-3-3784-8193. Fax:+81-3-3784-8247. E-mail: masaru-kato@umin.ac.jp.

Highlights

- Nanomedicines were captured by SPE discs after activation with 1 M metal solutions
- Bimodal pores of monolithic discs are suitable to enrich nanomedicines
- High recovery (nearly 100 %) of DOXIL from a 40× diluted solution was achieved

Download English Version:

<https://daneshyari.com/en/article/7607642>

Download Persian Version:

<https://daneshyari.com/article/7607642>

[Daneshyari.com](https://daneshyari.com)