

Accepted Manuscript

Title: Three-dimensional ionic liquid-ferrite functionalized graphene oxide nanocomposite for pipette-tip solid phase extraction of 16 polycyclic aromatic hydrocarbons in human blood sample

Authors: Yun Zhang, Yong-Gang Zhao, Wei-Sheng Chen, He-Li Cheng, Xiu-Qiong Zeng, Yan Zhu

PII: S0021-9673(18)30354-6
DOI: <https://doi.org/10.1016/j.chroma.2018.03.039>
Reference: CHROMA 359278

To appear in: *Journal of Chromatography A*

Received date: 29-1-2018
Revised date: 15-3-2018
Accepted date: 20-3-2018

Please cite this article as: Yun Zhang, Yong-Gang Zhao, Wei-Sheng Chen, He-Li Cheng, Xiu-Qiong Zeng, Yan Zhu, Three-dimensional ionic liquid-ferrite functionalized graphene oxide nanocomposite for pipette-tip solid phase extraction of 16 polycyclic aromatic hydrocarbons in human blood sample, *Journal of Chromatography A* <https://doi.org/10.1016/j.chroma.2018.03.039>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Three-dimensional ionic liquid-ferrite functionalized graphene oxide nanocomposite for pipette-tip solid phase extraction of 16 polycyclic aromatic hydrocarbons in human blood sample

Yun Zhang ^{a, 1}, Yong-Gang Zhao ^{a, b, 1}, Wei-Sheng Chen ^c, He-Li Cheng ^a, Xiu-Qiong

Zeng ^a, Yan Zhu ^{a*}

(^a Department of Chemistry, Zhejiang University, Hangzhou 310027, China; ^b Key Laboratory of Health Risk Appraisal for Trace Toxic Chemicals of Zhejiang Province, Ningbo Municipal Center for Disease Control and Prevention, Ningbo 315010, China;

^c Marine Environmental Monitoring Center of Ningbo, SOA, 315040, China)

*Corresponding author. E-mail address: zhuyan@zju.edu.cn (Y. Zhu).

¹These two are the first authors having given equal contribution.

***Highlights**

- PT-SPE method was developed for the enrichment of 16 PAHs in human blood.
- The PT-SPE procedure was solvent-saving, reusable, and needed less blood sample.
- The method provided referenced information for assessing risks of PAH exposure in China.

Download English Version:

<https://daneshyari.com/en/article/7608160>

Download Persian Version:

<https://daneshyari.com/article/7608160>

[Daneshyari.com](https://daneshyari.com)