

## Accepted Manuscript

How much separation for LC–MS/MS quantitative bioanalysis of drugs and metabolites?

Aimin Tan, John C. Fanaras



PII: S1570-0232(17)31658-6  
DOI: [doi:10.1016/j.jchromb.2018.03.019](https://doi.org/10.1016/j.jchromb.2018.03.019)  
Reference: CHROMB 21085

To appear in:

Received date: 25 September 2017  
Revised date: 3 January 2018  
Accepted date: 10 March 2018

Please cite this article as: Aimin Tan, John C. Fanaras , How much separation for LC–MS/MS quantitative bioanalysis of drugs and metabolites?. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Chromb(2017), doi:[10.1016/j.jchromb.2018.03.019](https://doi.org/10.1016/j.jchromb.2018.03.019)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**How much separation for LC–MS/MS quantitative bioanalysis of drugs and metabolites?**

Aimin Tan and John C. Fanaras

*(Nucro-Technics, Scarborough, ON, Canada)*

ACCEPTED MANUSCRIPT

---

*Note: This article was prepared based on the oral presentation given at the HPLC 2017 Prague.*

Download English Version:

<https://daneshyari.com/en/article/7615189>

Download Persian Version:

<https://daneshyari.com/article/7615189>

[Daneshyari.com](https://daneshyari.com)