


Accepted Manuscript

Myristic acid induces proteomic and secretomic changes associated with steatosis, cytoskeleton remodeling, endoplasmic reticulum stress, protein turnover and exosome release in HepG2 cells

Giulia Speziali, Laura Liesinger, Juergen Gindlhuber, Christina Leopold, Bettina Pucher, Jessica Brandi, Annalisa Castagna, Tamara Tomin, Petra Krenn, Gerhard G. Thallinger, Oliviero Olivieri, Nicola Martinelli, Dagmar Kratky, Matthias Schittmayer, Ruth Birner-Gruenberger, Daniela Cecconi



PII: S1874-3919(18)30167-2
DOI: doi:[10.1016/j.jprot.2018.04.008](https://doi.org/10.1016/j.jprot.2018.04.008)
Reference: JPROT 3104

To appear in: *Journal of Proteomics*

Received date: 9 October 2017
Revised date: 19 March 2018
Accepted date: 9 April 2018

Please cite this article as: Giulia Speziali, Laura Liesinger, Juergen Gindlhuber, Christina Leopold, Bettina Pucher, Jessica Brandi, Annalisa Castagna, Tamara Tomin, Petra Krenn, Gerhard G. Thallinger, Oliviero Olivieri, Nicola Martinelli, Dagmar Kratky, Matthias Schittmayer, Ruth Birner-Gruenberger, Daniela Cecconi, Myristic acid induces proteomic and secretomic changes associated with steatosis, cytoskeleton remodeling, endoplasmic reticulum stress, protein turnover and exosome release in HepG2 cells. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. *Jprot*(2017), doi:[10.1016/j.jprot.2018.04.008](https://doi.org/10.1016/j.jprot.2018.04.008)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Myristic acid induces proteomic and secretomic changes associated with steatosis, cytoskeleton remodeling, endoplasmic reticulum stress, protein turnover and exosome release in HepG2 cells

Giulia Speziali ¹, Laura Liesinger ^{2,3}, Juergen Gindlhuber ^{2,3}, Christina Leopold ⁴, Bettina Pucher ^{2,3,5}, Jessica Brandi ¹, Annalisa Castagna ⁶, Tamara Tomin ^{2,3}, Petra Krenn ^{2,3}, Gerhard G. Thallinger ^{3,5}, Oliviero Olivieri ⁶, Nicola Martinelli ⁶, Dagmar Kratky ^{4,7}, Matthias Schittmayer ^{2,3,4,7}, Ruth Birner-Gruenberger ^{2,3,4,7*}, Daniela Cecconi ^{1*}

¹ Department of Biotechnology, Proteomics and Mass Spectrometry Laboratory, University of Verona, Strada le Grazie 15, Verona, Italy

² Research Unit of Functional Proteomics and Metabolomics, Institute of Pathology, Medical University of Graz, Graz, Austria

³ Omics Center Graz, BioTechMed-Graz, Graz, Austria

⁴ Gottfried Schatz Research Center, Molecular Biology and Biochemistry, Medical University of Graz, Graz, Austria

⁵ Institute of Computational Biotechnology, Graz University of Technology, Graz, Austria

⁶ Department of Medicine, Unit of Internal Medicine, University of Verona, P.le L.A. Scuro 10, Verona, Italy

⁷ BioTechMed-Graz, Graz, Austria

* *Correspondence to:*

Prof. Daniela Cecconi, Department of Biotechnology, University of Verona

Strada le Grazie 15, 37134 Verona, Italy; FAX: +39 045 8027929; e-mail: daniela.cecconi@univr.it

Download English Version:

<https://daneshyari.com/en/article/7633437>

Download Persian Version:

<https://daneshyari.com/article/7633437>

[Daneshyari.com](https://daneshyari.com)