

## Author's Accepted Manuscript

Optimization of classification and regression analysis of four monoclonal antibodies from Raman spectra using collaborative machine learning approach

Laetitia Minh Mai Le, Balázs Kégl, Alexandre Gramfort, Camille Marini, David Nguyen, Mehdi Cherti, Sana Tfaily, Ali Tfayli, Arlette Baillet-Guffroy, Patrice Prognon, Pierre Chaminade, Eric Caudron



[www.elsevier.com/locate/talanta](http://www.elsevier.com/locate/talanta)

PII: S0039-9140(18)30227-3  
DOI: <https://doi.org/10.1016/j.talanta.2018.02.109>  
Reference: TAL18428

To appear in: *Talanta*

Received date: 25 October 2017  
Revised date: 24 February 2018  
Accepted date: 27 February 2018

Cite this article as: Laetitia Minh Mai Le, Balázs Kégl, Alexandre Gramfort, Camille Marini, David Nguyen, Mehdi Cherti, Sana Tfaily, Ali Tfayli, Arlette Baillet-Guffroy, Patrice Prognon, Pierre Chaminade and Eric Caudron, Optimization of classification and regression analysis of four monoclonal antibodies from Raman spectra using collaborative machine learning approach, *Talanta*, <https://doi.org/10.1016/j.talanta.2018.02.109>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Optimization of classification and regression analysis of four monoclonal antibodies from Raman spectra using collaborative machine learning approach

Laetitia Minh Mai Le<sup>a,b,1</sup> and Balázs Kégl<sup>c,g,1</sup>, Alexandre Gramfort<sup>c,e,f</sup>, Camille Marini<sup>c,d</sup>, David Nguyen<sup>a</sup>, Mehdi Cherti<sup>c,g</sup>, Sana Tfaili<sup>b,\*</sup>, Ali Tfayli<sup>b</sup>, Arlette Baillet-Guffroy<sup>b</sup>, Patrice Prognon<sup>a,b</sup>, Pierre Chaminade<sup>b</sup>, Eric Caudron<sup>a,b</sup>

<sup>a</sup> European Georges Pompidou Hospital (AP-HP), Pharmacy department, 75015 Paris, France

<sup>b</sup> Lip(Sys)<sup>2</sup>- EA7357 - Chimie Analytique Pharmaceutique (FKA EA4041 Groupe de Chimie Analytique de Paris-Sud), Univ. Paris-Sud, Université Paris-Saclay, F92290 Chatenay-Malabry, France.

<sup>c</sup> Center of Data Science, Université Paris-Saclay, 91440 Orsay, France.

<sup>d</sup> CMAP, Ecole Polytechnique, 91128 Palaiseau, France

<sup>e</sup> INRIA, Parietal team, Saclay, 91120 Palaiseau, France France

<sup>f</sup> LTCI, Télécom ParisTech, 75013 Paris, France

<sup>g</sup> LAL, CNRS, 91440 Orsay, France

**\*Corresponding author:** Sana Tfaili. 5, rue Jean Baptiste Clément, 92290 Châtenay-Malabry, France. sana.tfaili@u-psud.fr

## Abstract

The use of monoclonal antibodies (mAbs) constitutes one of the most important strategies to treat patients suffering from cancers such as hematological malignancies and solid tumors.

---

<sup>1</sup> Equally contributed to this article

Download English Version:

<https://daneshyari.com/en/article/7676576>

Download Persian Version:

<https://daneshyari.com/article/7676576>

[Daneshyari.com](https://daneshyari.com)