



ELSEVIER

Contents lists available at ScienceDirect

Talanta

journal homepage: www.elsevier.com/locate/talanta

Sensitive and selective determination of NO_2^- ion in aqueous samples using modified gold nanoparticle as a colorimetric probe



Yun-Sik Nam^{a,1}, Kwon-Chul Noh^{a,1}, Nak-Kyoon Kim^a, Yeonhee Lee^a, Hee-Kyung Park^b, Kang-Bong Lee^{a,*}

^a Advanced Analysis Center, Korea Institute of Science and Technology, P.O. Box 131, Cheongryangri, Seoul 130-650, Republic of Korea

^b Department of Oral Medicine and Oral Diagnosis, School of Dentistry, Seoul National University, 101 Daehak-ro Jongro-gu, Seoul 110-774, Republic of Korea

ARTICLE INFO

Article history:

Received 25 November 2013

Received in revised form

13 February 2014

Accepted 14 February 2014

Available online 22 February 2014

Keywords:

Nitrite ion

Gold nanoparticle

1-(2-mercaptoethyl)-1, 3, 5-Triazinane-2, 4,

6-trione

Colorimetric detection

ABSTRACT

A sensitive and selective colorimetric method for determination of nitrite ion in aqueous samples was developed using 1-(2-mercaptoethyl)-1, 3, 5-triazinane-2, 4, 6-trione-functionalized gold nanoparticles (MTT-GNPs). The nitrite ion seems to be used as a “molecular bridge”, which can form NH–N and NH–O hydrogen bonds with the MTT-GNPs, shorten the interparticle distance, and induce the aggregation of the MTT-GNPs. This aggregation results in a dramatic change from wine-red to purple-gray color. Therefore, the concentration of nitrite ion in environmental samples can be quantitatively detected using the MTT-GNPs sensor by the naked eyes or UV–vis spectrometer. Moreover, investigations have revealed the sensitivity of the detection could be clearly improved by modulating pH of the solution, which led to a more rapid color change in the optimized GNPs system. The absorption ratios (A_{790}/A_{535}) of the modified GNPs solution exhibited a linear correlation with nitrite ion concentrations and the limit of detection was 1 ppm. This cost effective sensing system allows for the rapid and facile determination of the concentration of NO_2^- ions in aqueous samples.

© 2014 Elsevier B.V. All rights reserved.

1. Introduction

When nitrite ions are used as preservatives for meat and fish products, they can react with secondary or tertiary amines present in the body resulting in the formation of carcinogenic nitrosamines, which enhances the possibility of cancer [1,2]. Furthermore, nitrite ions in the blood can react with iron(II) of hemoglobin forming methemoglobin, which lacks the ability to carry oxygen and causes fetal methemoglobin anemia, resulting in cyanosis [3]. Thus, the importance of the quantitative determination of nitrite ions in water, food, and agricultural products is widely recognized and significant, since a high concentration can lead to potential risks to human health. Specifically, the United States Environmental Protection Agency (US EPA) and World Health Organization (WHO) have established the maximum limit of nitrite ions in drinking water as 3 mg L^{-1} and 1 mg L^{-1} , respectively [3,4].

A variety of analytical methods have been reported for the determination of nitrite ions including electrometry [5,6],

chromatography [7–10], capillary electrophoresis [11], spectrophotometry [12,13], and spectrofluorimetry [14], but most of these methods require complicated instruments and professional staff. Hence, exploring an accurate, rapid, and convenient analytical method to detect nitrite ions still remains a challenge for analytical chemists.

With developments in nanotechnology, various novel colorimetric assays have been exploited. Gold nanoparticles (GNPs) have been extensively used as colorimetric probes for different sensing and biosensing applications owing to their unique size-dependent and interparticle-distance dependent optical properties and accompanying color changes. GNP-based colorimetric methods have been reported for the detection of anions [15,16], heavy metal ions [17–21], DNA [22], bacteria [23], and cancerous cells [24], among others. Lu's group [25] has developed 1-(2-mercaptoethyl)-1,3,5-triazinane-2,4,6-trione (MTT)-stabilized GNPs as a colorimetric probe for melamine detection in milk, based on the hydrogen-bonding between modifiers and melamine. Therefore, the development of on-site, real-time, and low-cost melamine sensing has become increasingly attractive. Furthermore, MTT can be easily conjugated to GNPs through the –SH group. Inspired by these works, we have examined the potential of MTT-modified gold nanoparticles as a novel and sensitive probe for the

* Corresponding author. Tel.: +82 2 958 5957; fax: +82 2 958 5969.

E-mail address: leekb@kist.re.kr (K.-B. Lee).

¹ These authors contributed equally to this work.

determination of negatively charged ions in aqueous samples, since the NH of the MTT molecule could bind anions via hydrogen bonding. Here, we have tested the MTT-GNPs with various anions, which are F^- , Cl^- , Br^- , SO_4^{2-} , PO_4^{3-} , NO_3^- , and NO_2^- . The NH of MTT-GNPs could only be cross-linked with nitrite ions, accompanying clear color change (from wine-red to purple gray). Moreover, the GNP-based system has a better sensitivity for the detection of nitrite ions in acidic solution. Also, the absorption ratio (A_{790}/A_{535}) of the modified GNPs in the pH-optimized system exhibits a linear correlation with the nitrite concentration, allowing the determination of the nitrate concentration in aqueous samples within 30 min. Therefore, an optimized MTT-GNP system as a novel, on-site, real-time, and low-cost nitrite ion sensor has a wide range of practical applications.

2. Experimental

2.1. Materials

Gold(III) chloride hydrate, sodium citrate, ethyl chloroformate, acetone, and potassium thiocyanate were purchased from Sigma-Aldrich (St. Louis, MO, USA). Methylene chloride and HCl were purchased from J. T. Baker (Phillipsburg, NJ, USA). F^- , Cl^- , Br^- , SO_4^{2-} , PO_4^{3-} , NO_3^- , and NO_2^- were purchased from AccuStandard (New Haven, CT, USA). pH paper was purchased from Whatman International Ltd. (Maidstone, UK). HCl and NaOH were purchased from Samchun Chemical (Gyeong Gi-Do, Korea). Sodium chloride was purchased from Junsei Chemical (Tokyo, Japan). Distilled water used in this experiment was obtained through a Milli-Q water purification system from Millipore (Bedford, MA, USA) and was used throughout the study.

2.2. Apparatus

The absorption spectra of the GNPs were recorded at room temperature using a UV–vis–NIR spectrophotometer 5000 (Varian Inc., Palo Alto, CA, USA). UV–vis was measured in the range of 400–800 nm using quartz cells of 1 mm path length. Measurement of particle size distributions were carried out using a Zetasizer (Malvern Instruments Ltd., Worcestershire, UK). The image and the diameter of the MTT-GNP and the nitrite-induced aggregation of MTT-GNP were measured from a micrograph using a transmission electron microscope (TEM) CM30 (Philips, NC, USA) and an atomic force microscope (AFM) XFM XE-100 (Park systems, Suwon, Republic of Korea). TEM specimens were obtained by depositing a dispersion of the GNPs and evaporating the solvent. The concentrations of nitrite ion in aqueous solutions were measured using ion chromatography (IC) (Metrohm, Riverview, FL, USA).

2.3. Preparation of MTT and GNPs conjugated with MTT

MTT was synthesized according to literature procedure [26], and then MTT-conjugated gold nanoparticles were prepared by a ligand-exchange reaction between MTT and citrate-stabilized gold nanoparticles. We synthesized ~33 nm gold nanoparticles by the reduction of $HAuCl_4$ using sodium citrate. Generally, 50 mL of 1 mM $HAuCl_4$ as heated to reflux with stirring, and then 5 mL of 38.8 mM sodium citrate was added rapidly. The solution was boiled for another 30 min to give a wine-red colored solution. The ligand-exchange reaction was performed at room temperature by mixing 0.5 mL of the prepared gold colloids with 1 mL of aqueous 1 mM MTT under stirring. Volumes of 0.08, 0.3, 0.5, 0.7, 0.9, and 1.1 mL of 1 mM MTT were added to 0.5 mL of GNPs. We did not observe any color change for the MTT-GNP solution upon

addition of MTT, indicating that the self-aggregation of MTT-GNPs did not occur.

3. Results and discussion

Nanomaterials often display unique physical properties that are not observed in the same materials on a larger scale. The coherent collective motion of electrons on the surface in noble metals such as gold or silver arises when the size of the particles are on the order of a few nanometers. Gold nanoparticles strongly absorb in the visible region of light, when the frequency of the incident electromagnetic field is resonant with the coherent oscillation of the electrons on the surface of the nanoparticle. In nanoparticles, this phenomenon is called surface plasmon resonance (SPR). A strong absorption peak of SPR appears at about 520 nm in the UV–vis spectra of gold nanoparticles. This is the reason why gold nanoparticles display a wine-red color in solution. The SPR peak frequency is influenced by the size and shape of the nanoparticles, the refractive index of the surrounding medium, and the degree of aggregation [27–29].

We synthesized MTT, a thiol-functionalized cyanuric acid [26], and prepared ~33 nm MTT-GNPs using the ligand-exchange reaction of GNPs with MTT [25]. The GNP size under these preparation conditions is larger than that (12 nm) of MTT-stabilized GNPs synthesized in an earlier study [25]. The mean size of the GNPs was easily affected by the amount of sodium citrate and other preparation conditions. It is known that the GNP size affects the surface plasma absorption maxima. The λ max values of GNPs of sizes 15 and 33 nm were 519 and 535 nm, respectively. However, this change in λ max as a function of GNP size seems not to influence the results and reliability of the experiments [27–29]. The gold nanoparticles conjugated with MTT were well dispersed and displayed uniform wine-red color due to the strong SPR at 535 nm.

Following the synthesis of the MTT-GNPs, we developed a simple colorimetric assay for the selective detection of anions. The control solution that did not contain any anions showed the wine-red color, and each solution with the addition of F^- , Cl^- , Br^- , PO_4^{3-} , SO_4^{2-} or NO_3^- ions (5 ppm) did not show any color change. However, the solution with nitrite ion gradually changed from a wine-red to a purple-gray color as displayed in Fig. 1(A). The UV–vis absorption spectra of the MTT-GNPs solutions were acquired 30 min. after the addition of each anion, and are displayed in Fig. 1 (B). λ_{790} does not seem to be λ max as shown in Fig. 1(B), and λ max could be a little longer than 800 nm. The reason why we chose A_{790} as λ max is because the radiation source of the UV–vis spectrometer used in this experiment only covers the range 200–800 nm, and the absorbance curves of the UV–vis spectra drop vertically at 800 nm. But we are certain that this choice does not make any difference to the relative absorbance ratios of the two λ max values.

The sensitivity and selectivity of GNPs towards various anions was further studied by plotting the absorbance ratios (A_{790}/A_{535}) of the MTT-GNPs solution in the presence of each anion as shown in Fig. 1(C).

The selectivity of the optimized sensor for nitrite ions was evaluated by comparison of the absorbance ratio (A_{790}/A_{535}) with that of solutions containing similar anions (F^- , Cl^- , Br^- , PO_4^{3-} , SO_4^{2-} , and NO_3^-). It was found that GNPs conjugated with MTT responded selectively to nitrite, as indicated by the dramatic increase in the absorbance ratio (A_{790}/A_{535}). Thus, we successfully identified the aggregation of nanoparticles by nitrite ion in GNPs conjugated with MTT. The absorbance at 535 nm decreased and a new band with an absorption at 790 nm developed, and could be attributed to the coupled plasmon absorbance by closer contact

Download English Version:

<https://daneshyari.com/en/article/7680271>

Download Persian Version:

<https://daneshyari.com/article/7680271>

[Daneshyari.com](https://daneshyari.com)