Accepted Manuscript

Title: Current approaches in sample preparation for trace analysis of selected endocrine disrupting compounds. focus on polychlorinated biphenyls, alkylphenols and parabens.

Author: Tomasz Grześkowiak, Beata Czarczyńska-Goślińska, Agnieszka Zgoła-Grześkowiak

 PII:
 S0165-9936(15)00250-2

 DOI:
 http://dx.doi.org/doi:10.1016/j.trac.2015.07.005

 Reference:
 TRAC 14523

To appear in: Trends in Analytical Chemistry



Please cite this article as: Tomasz Grześkowiak, Beata Czarczyńska-Goślińska, Agnieszka Zgoła-Grześkowiak, Current approaches in sample preparation for trace analysis of selected endocrine disrupting compounds. focus on polychlorinated biphenyls, alkylphenols and parabens., *Trends in Analytical Chemistry* (2015), http://dx.doi.org/doi:10.1016/j.trac.2015.07.005.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

1	Current approaches in sample preparation for trace analysis of selected
2	endocrine disrupting compounds. Focus on polychlorinated biphenyls,
3	alkylphenols and parabens.
4	
5	Tomasz Grześkowiak ¹ , Beata Czarczyńska-Goślińska ² , Agnieszka Zgoła-Grześkowiak ^{1*}
6	
7	¹ Institute of Chemistry and Technical Electrochemistry, Poznan University of Technology,
8	Berdychowo 4, 60-965 Poznań, Poland,
9	² Department of Pharmaceutical Technology, Poznan University of Medical Sciences,
10	Grunwaldzka 6, 60-780 Poznań, Poland
	S
11	
12	Highlights
13	Sample preparation methods for isolation of endocrine disruptors were presented
14	Methods for polychlorinated biphenyls, alkylphenols and parabens are included
15	Ultrasonic assisted and pressurised liquid extraction dominate for solid matrix
16	Solid-phase extraction and microextraction dominate for liquid matrix
17	Trends include new sorbents, simplification and green chemistry approach
18	
	~ G
19	Abstract
20	Endocrine disrupting compounds (EDC) belong to the most widely analyzed chemicals.
21	This review covers three different groups of these compounds: polychlorinated biphenyls,
22	alkylphenols and parabens with the last group belonging to suspected EDC. Current trends in
23	sample preparation of these compounds include usage of different techniques aiming mainly
24	at simplification of procedures and low solvent consumption. Pressurised liquid extraction
25	and ultrasonic assisted extraction are the main techniques used for isolation of EDC from the

Download English Version:

https://daneshyari.com/en/article/7689158

Download Persian Version:

https://daneshyari.com/article/7689158

Daneshyari.com