

Author's Accepted Manuscript

Effect of particle size and local disorder on specific surface area of layered double hydroxides upon calcination-reconstruction

Bo-Kyung Kim, Gyeong-Hyeon Gwak, Tomohiko Okada, Jae-Min Oh



PII: S0022-4596(18)30130-0
DOI: <https://doi.org/10.1016/j.jssc.2018.03.041>
Reference: YJSSC20167

To appear in: *Journal of Solid State Chemistry*

Received date: 13 December 2017
Revised date: 29 March 2018
Accepted date: 31 March 2018

Cite this article as: Bo-Kyung Kim, Gyeong-Hyeon Gwak, Tomohiko Okada and Jae-Min Oh, Effect of particle size and local disorder on specific surface area of layered double hydroxides upon calcination-reconstruction, *Journal of Solid State Chemistry*, <https://doi.org/10.1016/j.jssc.2018.03.041>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Effect of particle size and local disorder on specific surface area of layered double hydroxides upon calcination-reconstruction

Bo-Kyung Kim¹, Gyeong-Hyeon Gwak², Tomohiko Okada^{3*}, Jae-Min Oh^{1*}

¹Department of Chemistry and Medical Chemistry, College of Science and Technology, Yonsei University, Wonju, Gangwondo, 26493, Republic of Korea

²Beamline Research Division, Pohang Accelerator Laboratory, Pohang University of Science and Technology, Pohang, Gyeongsangbukdo, 37673, Republic of Korea

³Department of Chemistry and Material Engineering, Faculty of Engineering, Shinshu University, Wakasato 4-17-1, Nagano 380-8553, Japan

Correspondence

*Tel. +82-33-760-2368; fax: +82-33-760-2182; jaemin.oh@yonsei.ac.kr , ORCID: 0000-0003-1638-9957

*Tel. +81-26-269-5414; fax: +81-26-269-5424; tomohiko@shinshu-u.ac.jp ORCID: 0000-0002-9361-7004

Abstract

The specific surface area changes in MgAl-layered double hydroxide (LDH), which is also known as hydrotalcite, were monitored along with their changes in particle size, crystallinity

Download English Version:

<https://daneshyari.com/en/article/7757679>

Download Persian Version:

<https://daneshyari.com/article/7757679>

[Daneshyari.com](https://daneshyari.com)