



Design and synthesis of plant cyclopeptide Astin C analogues and investigation of their immunosuppressive activity



Fei Li^{a,c,d}, Xi-Xi Guo^{b,d}, Guang-Zhi Zeng^a, Wei-Wei Qin^b, Bo Zhang^b, Ning-Hua Tan^{a,b,*}

^a State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, People's Republic of China

^b State Key Laboratory of Natural Medicines, Department of TCMs Pharmaceuticals, School of Traditional Chinese Pharmacy, China Pharmaceutical University, Nanjing 211198, People's Republic of China

^c University of Chinese Academy of Sciences, Beijing 10049, People's Republic of China

ARTICLE INFO

Keywords:

Astin C analogues
Solid-phase peptide synthesis
Immunosuppressive activity

ABSTRACT

To further investigate on the structure-activity relationships of immunosuppressive Astin C, seventeen analogues **1–17** were designed and synthesized *via* amino acid substitution strategy by the solid-phase peptide synthesis method for the first time. In comparison with Astin C ($IC_{50} = 12.6 \pm 3.3 \mu M$), only compounds **2** ($IC_{50} = 38.4 \pm 16.2 \mu M$), **4** ($IC_{50} = 51.8 \pm 12.7 \mu M$), **5** ($IC_{50} = 65.2 \pm 15.6 \mu M$), and **8** ($IC_{50} = 61.8 \pm 12.4 \mu M$) exhibited immunosuppressive activity in the Lymph node cells of mice. These results showed that the Astin C analogues containing D -amino acid residues, hydrophobic long-chain alkyl substituents, and aryl substituents performed better than those carrying hydrophilic amino acid residues and short-chain alkyl substituents. Moreover compounds **15**, **16**, and **17** had no immunosuppressive activity, which suggested that *cis*-3,4-dichlorinated proline played an important role in the immunosuppressive activity of Astin C.

In our project on unique structure and potential activity of cyclopeptides from traditional Chinese medicines (TCMs), we have found that Astins, a family of homomonocyclopeptides, named as Compositae-type cyclopeptides, were only isolated from the roots and rhizomes of *Aster tataricus* L. (Compositae)^{1–4}, which is one of TCMs used for relieving cough and eliminating phlegm.⁵ It is noteworthy that Astins share an unusual structural similarity, which are mainly characterized by the presence of four unnatural amino acids (L -Abu, L -allo-Thr, β -Phe, and mono- or di-chlorinated L -Pro residues) and one proteinogenic amino acid (L -Ser).¹ So far, sixteen cyclopentapeptides (Astins A–I¹, K–P²) together with two unique skeleton cyclopeptides, Tataricins A and B³ (Fig. 1), have been isolated from *A. tataricus*. Previous studies have shown that Astin A, B, and C with the *cis*-3,4-dichlorinated proline residue played an important role in antitumor activity.^{6,7} Cozzolino et al. have reported that synthesized antineoplastic cyclic astin analogues can kill tumor cells *via* caspase-mediated induction of apoptosis.⁸ Recently we have demonstrated that Astin C could induce mitochondria-dependent apoptosis of activated T cells, and exhibited potential immunosuppressive activity *in vitro* and *in vivo*.⁴ To the best of our knowledge, Astin C and its derivatives with a chlorinated proline have not been synthesized. There are only two reports about the synthesis of

Astins. The first is Astin G, which was totally synthesized in 1999.⁹ The second is Tataricins A and B, which were synthesized by our group in 2013.³ However, study on the structure-immunosuppressive activity of Astin C has not been carried out to date. Herein, several amino acid substitution strategies have been conducted to modify Astin C and seventeen analogues were synthesized for the first time by the solid-phase peptide synthesis (SPPS) method. Their immunosuppressive activities were screened against Lymph node cells and the structure-activity relationships (SARs) have been discussed.

Astin C contains a cyclic pentapeptide core comprised of two L -Abu, one L -Ser, one β -Phe, and one strikingly *cis*-3,4-dichlorinated L -Pro. Its structure was divided into two parts, *i.e.* substitution part and conserved sequence part (Fig. 2). Firstly, we synthesized two analogues by the SPPS method, which bear 3,4-dehydro L -Pro in compound **15** and *cis*-3,4-dihydroxylated L -Pro in compound **16**, to study the role of the *cis*-3,4-dichlorinated L -Pro on the immunosuppressive activity against lymph node cells. Secondly, we synthesized other fifteen Astin C analogues **1–14** and **17** *via* amino acid substitution strategy by the SPPS method to study their SARs.

Firstly, we prepared the intermediate *cis*-3,4-diol- L -Pro¹⁰ **P-5a**, and the synthetic route is shown in Scheme 1. Compound **P-1** was prepared

* Corresponding author at: State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, People's Republic of China.

E-mail address: nhtan@cpu.edu.cn (N.-H. Tan).

^d These authors contributed equally to this work.




Fig. 1. Structures of Astins family.





Fig. 2. Substitution of the structure of Astin C.

starting with commercially available *trans*-4-hydroxy-L-proline and Boc₂O in the presence of NaHCO₃ in dioxane/H₂O, which reacted with BnBr in DMF to give compound **P-2** in excellent yield. By using Appel-Lee reaction, compound **P-2** was successfully converted to compound **P-3** in 97% yield. Subsequently, elimination of compound **P-3** with DBU in ACN afforded compound **P-4a** as a major product, which is oxidized by K₂OsO₂(OH)₄ to the diol **P-5a** as the major product.

With 3,4-dehydro-L-Pro **P-4a** and *cis*-3,4-diol-L-Pro **P-5a** in hand, *cis*-3,4-dichlorinated L-Pro in Astin C was then modified. As depicted in Scheme 2, *cis*-3,4-dichlorinated L-Pro was substituted by Ala (Ala scan¹¹), **P-4a**, and **P-5a**, respectively, and compounds **14**, **15**, and **16** were obtained by the SPPS method similar to Scheme 3. In addition, cyclotetrapeptide **17** was prepared by cutting the *cis*-3,4-dichlorinated proline residue.



Scheme 1. Reagents and conditions: (a) Boc₂O, NaHCO₃, dioxane : H₂O = 1 : 1, rt, 24 h, 86%; (b) BnBr, K₂CO₃, DMF, rt, overnight, 92%; (c) Ph₃P, I₂, imidazole, DCM, N₂, reflux, 12 h, 97%; (d) DBU, ACN, reflux, overnight, 78%; (e) K₂OsO₂(OH)₄, K₃Fe(CN)₆, K₂CO₃, *t*-BuOH : H₂O = 1 : 1, rt, 24 h, 83%.

Next, solid-phase synthesis of other Astin C analogues were carried out by substituting the *cis*-3,4-dichlorinated proline residue with various amino acids. Considering that Astins is all consisted of L-amino acid, we firstly introduced D-Thr and D-*allo*-Thr to synthesize Astin C analogues **2** and **4**. Moreover, hydrophobic amino acid residues (L-Ada, L-Val, L-Leu, β-Phe, L-Tyr) and hydrophilic amino acid residues (L-Thr, L-*allo*-Thr, L-Lys, L-Arg, L-His, (2S, 3S)-3-O-(pyrrole-2-carboxyloxy)-*allo*-Thr) were also introduced to prepare Astin C analogues **1**, **3**, and **5–13**, respectively. Because of the same sequence of linear tetrapeptide with L-Abu⁵-β-Phe⁴-L-Ser³-L-Abu² in **1–16**, we designed to cyclize at the Abu⁵/substituted amino acid¹ site. As an example, the synthetic route of compound **2** was shown in Scheme 3.

Compound **2** was synthesized using the Fmoc-Abu-CTC Resin. Fmoc group protecting amino acids were successively coupled by HBTU and DIPEA in DMF. 20% piperidine/DMF was used for deblocking. The reaction was monitored by ninhydrin color reaction. After the last step, the peptide resin was washed with DMF (three times), and MeOH (three times), then dried by N₂ bubbling over night. TFA in DCM was added to the flask containing the peptide resin to release the crude linear peptide from the resin. Cyclization was accomplished by using HATU and DIPEA. The progress of the cyclization was monitored by LC-MS. The solvent was removed under reduced pressure and the crude peptide was purified by Prep-HPLC to get the final product **2**.¹² Similarly, compounds **1**, **3–13** were prepared by the above synthetic method.

With Astin C and seventeen analogues **1–17** in hand (Fig. 3), we have screened immunosuppressive activity against lymph node cells, in which CsA was used as the positive control. As shown in Table 1, Astin C shows potential immunosuppressive activity. Compared with Astin C (IC₅₀ = 12.6 ± 3.3 μM), compound **2** (IC₅₀ = 38.4 ± 16.2 μM) with D-Thr residue and compound **4** (IC₅₀ = 51.8 ± 12.7 μM) with D-*allo*-Thr residue showed comparable immunomodulatory activity, which opened doors to us that the introduction of D-amino acids is a moderate strategy. On the other hand, compound **5** (IC₅₀ = 65.2 ± 15.6 μM) with hydrophobic long-chain alkyl substituent and compound **8** (IC₅₀ = 61.8 ± 12.4 μM) with β-Phe residue (aryl substituent) also exhibit comparable immunosuppressive activity. Unfortunately, we

Download English Version:

<https://daneshyari.com/en/article/7778269>

Download Persian Version:

<https://daneshyari.com/article/7778269>

[Daneshyari.com](https://daneshyari.com)