



Contents lists available at ScienceDirect

Bioorganic & Medicinal Chemistry Letters

journal homepage: www.elsevier.com/locate/bmcl

Design and synthesis of novel dasatinib derivatives as inhibitors of leukemia stem cells

Hui-ying Li^{b,c}, Ding-Di He^{a,b,c}, Xiu-Juan Zhao^{a,b}, Tong-Yan Sun^{a,b}, Quan Zhang^a, Cui-Gai Bai^{b,*}, Yue Chen^{a,*}

^aThe State Key Laboratory of Medicinal Chemical Biology, Collaborative Innovation Center of Chemical Science and Engineering (Tianjin), College of Pharmacy, and Tianjin Key Laboratory of Molecular Drug Research, Nankai University, Tianjin 300353, PR China

^bHigh-throughput Molecular Drug Discovery Center, Tianjin International Joint Academy of BioMedicine, Tianjin 300457, PR China

ARTICLE INFO

Article history:

Received 22 November 2017

Revised 5 January 2018

Accepted 11 January 2018

Available online 12 January 2018

Keywords:

Dasatinib derivatives

Biososteres

Triazole

Oxadiazole

Colony formation

Cell migration

ABSTRACT

We used the concept of biososteres to design and synthesize a novel series of dasatinib derivatives for the treatment of leukemia. Unfortunately, most of the dasatinib derivatives did not show appreciable inhibition against leukemia cell lines K562 and HL60. However, acrylamide compound **2c** had comparable inhibitory activity with dasatinib against K562 cells ($IC_{50} = 0.039$ nM vs. 0.069 nM). And amide compound **2a** and acrylamide compound **2c** also had comparable inhibitory activity with dasatinib against the leukemia cell line HL60 ($IC_{50} = 0.25$ nM and 0.26 nM vs. 0.11 nM). Against the leukemia progenitor cell line KG1a, triazole compounds **15a** and **15d–15f** and oxadiazole compounds **24a–24d** were more potent than dasatinib. In particular, the hydroxyl compounds **15a** and **24a** were about 64 and 180 fold more potent than dasatinib against KG1a cells ($IC_{50} = 0.14$ μ M and 0.05 μ M vs. 8.98 μ M). Compounds **15a** and **24a** also inhibited colony formation in MCF-7 cells and inhibited cell migration in the cell wound scratch assay in B16BL6 cells. Moreover, hydroxyl compounds **15a** and **24a** had low toxicity *in vivo*.

© 2018 Elsevier Ltd. All rights reserved.

Dasatinib is a second-generation, orally available, potent, and multi-targeted inhibitor of Bcr-Abl and Src family kinases.^{1,2} Dasatinib is effective in imatinib-resistant wild type and mutant BCR-ABL cell lines, except in those carrying the T315I mutation.^{3–7} Dasatinib also has high selectivity for normal hematopoietic cells and leukemia cells.⁸ When dasatinib was used in conjunction with bone marrow transplantation, however, many mutations emerged, including T315I/A, F317L/I/C/V, V299L ($IC_{50} > 15$ nM), Y253F, E255 K/V and G250E.^{9–14} Dasatinib also fails to eliminate the quiescent chronic myeloid leukemia (CML) stem cell compartment.¹⁵

Biososterism is a very important strategy for drug design in medicinal chemistry, and 1,3,4-oxadiazole and 1,2,3-triazole are considered to be biososteres of the amide group. 1,3,4-Oxadiazole and 1,2,3-triazole rings have special biological properties and these moieties have been successfully used in multi-targeted drugs, including the anti-cancer drug, zibotentan.¹⁶

Here, a series of dasatinib derivatives, containing oxadiazole and triazole rings, was synthesized and the activity of the com-

pounds was evaluated in the leukemia cell lines K562 and HL60, and the leukemia progenitor cell line, KG1a. Lead compounds were further evaluated by the colony forming assay in MCF-7 cells and the cell wound scratch assay in B16BL6 cells. Preliminary toxicity was also discussed.

Dasatinib derivatives were prepared using the sequences shown in Schemes 1–5.


Firstly, acid amide compounds **2a–2d** were synthesized (Scheme 1). Dasatinib was treated with phthalimide to provide compound **1**, which was deprotected to provide amine compound **2a**. Amine compound **2a** was then converted to compounds **2b–2d**. Treatment with CS₂ under basic conditions afforded isothiocyanate compound **2b**, treatment with acryloyl chloride provided the allyl amide compound **2c** and treatment with chloroacetyl chloride gave the chloroethyl amide compound **2d**.

Triazole ring compounds **15a–15f** were synthesized as shown in Schemes 2 and 3. Boc protection of amine compound **3** provided compound **4**, which was converted to iodide compound **5** using *N*-iodosuccinimide (NIS). Iodide compound **5** was reacted with trimethylsilylacetylene to provide compound **6**, which was then deprotected to give alkyne compound **7**. Azide compound **9** and alkyne compound **7** were treated with Cu₂O nanoparticles to give the triazole ring compound **10** using click chemistry. The Boc

* Corresponding authors.

E-mail addresses: baicuigai@tjlab.org (C.-G. Bai), yuechen@nankai.edu.cn (Y. Chen).

^c These authors contributed equally to this work.



Scheme 1. Reagents and conditions: (a) PPh_3 , phthalimide, DIAD, THF, r.t., 88%; (b) $\text{N}_2\text{H}_2\cdot\text{H}_2\text{O}$, EtOH, 78 °C, 72%; (c) DCC, CS_2 , CH_2Cl_2 , r.t., 52%; (d) acryloyl chloride, Et_3N , CH_2Cl_2 , r.t., 49%; (e) chloroacetyl chloride, Et_3N , CH_2Cl_2 , r.t., 55%;

protecting group was then removed from compound **10** to provide compound **11**. Condensation of compounds **11** and **12** gave compound **13**, which was treated with different amines to provide compounds **15a** and **15b**. Triazole amine compound **15c** was prepared from compound **15b** by removal of the Boc group. Compounds **15d–15f** were synthesized using the same procedures for synthesis of acid amide compounds **2b–2d**, respectively.

1,3,4-Oxadiazole compounds **24a–24e** were prepared as shown in Schemes 4 and 5. Amine compound **16** was treated with compound **12** to afford pyrimidine **17**, which was condensed with different amine by condensation reaction to provide compounds **19a** and **19b**. Compounds **19a** and **19b** were treated with hydrazine solution to provide carbonylhydrazone compounds **20a** and **20b**, which were treated with chloride **21** to give compounds **22a** and **22b**. Compounds **22a** and **22b** were then converted to compounds **23a** and **23b**. Deprotection of compounds **23a** and **23b** gave compounds **24a** and **24b**. Compound **24b** was then converted to compounds **24c–24e**, using the procedures described for the synthesis of compounds **2b–2d**.

Most of the dasatinib derivatives did not noticeably inhibit the growth of K562 and HL60 cells, but both triazole and oxadiazole compounds were more potent than dasatinib against leukemia progenitor cell line KG1a (Table 1).

For compounds **2a–2d**, inhibitory activities did not increase. In K562 cells, the IC_{50} values of compounds **2a–2d** were 0.72, 3.77, 0.039 and 1.67 nM, respectively. The inhibitory activity of acrylamide compound **2c** was comparable to that of dasatinib (IC_{50} = 0.069 nM). In HL60 cells, the activities of compounds **2a** and **2c** were very similar to that of dasatinib (IC_{50} = 0.25 nM and 0.26 vs. 0.11 nM). Isothiocyanate compound **2b** (IC_{50} = 1.07 nM) had comparable inhibitory activity to that of compound **2d** (IC_{50} = 0.86 nM). In KG1a cells, isothiocyanate compound **2b** (IC_{50} = 1.96 μM)

was more potent than dasatinib (IC_{50} = 8.98 μM) whereas compounds **2a**, **2c** and **2d** (IC_{50} = 25.39, 12.88, and 13.56 μM , respectively) were less potent than dasatinib.

Inhibitory activities of triazoles **15a** and **15c–15f** were reduced 170–530-fold against K562 cells and 35–150-fold against HL60 cells whereas inhibitory activities were enhanced 4–64-fold against leukemia progenitor cells KG1a. Amongst the triazoles, isothiocyanate compound **15d** had the lowest inhibitory activity against K562 and HL60 cells (IC_{50} = 36.48 and 16.57 nM, respectively) and compounds **15a**, **15c**, **15e** and **15f** had comparable inhibitory activities (IC_{50} = 14.14, 11.73, 15.09 and 17.65 nM, respectively, against K562 cells and 4.19, 4.08, 5.46 and 9.82 nM, respectively, against HL60 cells). In KG1a cells, amine compound **15c** had the poorest inhibitory activity (IC_{50} = 2.06 μM) and compound **15a** had best inhibitory activity (IC_{50} = 0.14 μM). Compounds **15d–15f** had comparable inhibitory activities (IC_{50} = 0.32, 0.25 and 0.45 μM).

The IC_{50} values of the oxadiazole compounds against both K562 and HL60 cell lines were at least an order of magnitude higher than that of dasatinib. Against KG1a cells, compound **24e** (IC_{50} = 9.75 μM) had comparable inhibitory activity to dasatinib (IC_{50} = 8.98 μM). Compounds **24a–24d** (IC_{50} = 0.05, 0.68, 0.15, and 0.11 μM , respectively) were more potent than dasatinib. It is noteworthy that compound **24a** was about 180-fold more potent than dasatinib.

Dasatinib, and compounds **15a** and **24a**, significantly and dose-dependently decreased the formation of colonies in MCF-7 cells, compared with the control group (Fig. 1). The high dose of compound **15a** was more potent than dasatinib and the high dose of compound **24a** was comparable with dasatinib. Compounds **15a** and **24a** could thus inhibit the growth of cancer stem cells, and compound **15a** was more potent than compound **24a**.

Download English Version:

<https://daneshyari.com/en/article/7779658>

Download Persian Version:

<https://daneshyari.com/article/7779658>

[Daneshyari.com](https://daneshyari.com)