



## Dodecyl sorbitan ethers as antimicrobials against Gram-positive bacteria



Dorine Belmessieri<sup>a,b,c</sup>, Charlotte Gozlan<sup>a,d</sup>, Marie-Christine Duclos<sup>a</sup>, Oana Dumitrescu<sup>b,c</sup>,  
G rard Lina<sup>b,c,\*</sup>, Andreas Redl<sup>d</sup>, Nicolas Duguet<sup>a,\*</sup>, Marc Lemaire<sup>a,\*</sup>

<sup>a</sup> Univ Lyon, Universit  Claude Bernard Lyon1, CNRS, INSA-Lyon, CPE-Lyon, Institut de Chimie et Biochimie Mol culaires et Supramol culaires, (ICBMS), UMR 5246, Equipe CAtalyse, SYnth se et ENvironnement (CASYEN), 43 boulevard du 11 novembre 1918, F-69622 Villeurbanne Cedex, France

<sup>b</sup> Univ Lyon, Universit  Claude Bernard Lyon 1, Ecole Normale Sup rieure de Lyon, CNRS, UMR5308, International Center for Infectiology Research (CIRI), Inserm U1111, 7 rue Guillaume Paradin, F-69008 Lyon, France

<sup>c</sup> Centre National de R f rence des Staphylocoques, 59 boulevard Louis Pinel, F-69677 Bron Cedex, France

<sup>d</sup> Tereos Syral SAS, Z.I et Portuaire, B.P.32, 67390 Marckolsheim, France

## ARTICLE INFO

## Article history:

Received 19 July 2017

Revised 5 September 2017

Accepted 6 September 2017

Available online 7 September 2017

## Keywords:

Acetals

Antimicrobials

Ethers

Resistant bacteria

Sorbitan

## ABSTRACT

A range of amphiphilic sorbitan ethers has been synthesized in two steps from sorbitan following an acetalization/hydrogenolysis sequence. These sorbitan ethers and the acetal intermediates have been evaluated as antimicrobials against Gram-negative and Gram-positive bacteria. No antimicrobial activity was observed for Gram-negative bacteria. However, the compounds bearing a linear dodecyl chain exhibit antimicrobial activity (MIC as low as 8  $\mu\text{g/mL}$ ) against Gram-positive bacteria such as *Listeria monocytogenes*, *Enterococcus faecalis* and *Staphylococcus aureus*. Encouraged by these preliminary results, dodecyl sorbitan was tested against a range of resistant strains and was found to be active against vancomycin-, methicillin- and daptomycin-resistant strains (MIC = 32–64  $\mu\text{g/mL}$ ).

  2017 Elsevier Ltd. All rights reserved.

Antimicrobials are essential compounds to prevent and cure bacterial infections.<sup>1</sup> In this respect, they are widely used in medicine but also in other fields such as in cosmetics, food,<sup>2</sup> textile<sup>3</sup> and packaging.<sup>4</sup> However, the misuse and overuse of antimicrobials have led to the emergence of antimicrobial resistance (AMR). AMR presents a serious problem for public health as it threatens the prevention and the treatment of bacterial infections and compromises the success of major surgery and cancer chemotherapy.<sup>5</sup> More worrying, humanity is now facing infections by superbugs that are resistant to all known antimicrobial chemotherapy.<sup>6</sup> In this context, the development of new effective antimicrobials is highly desirable.


Sorbitan esters, known under the trade name Span<sup> </sup>, are surface active compounds with very low impacts on health (i.e. no acute toxicity) and environment (i.e. readily biodegradable).<sup>7</sup> For example, sorbitan monolaurate (Span 20, E493), one of the most representative compound of this family, is widely used in food,<sup>8</sup> pharmaceutical and cosmetic industries.<sup>9</sup> Similarly to other carbohydrate fatty esters such as sucrose monolaurate,<sup>10</sup> lauroylmaltose and maltotriose,<sup>11</sup> and methyl 6-O-lauroyl- -D-glucopyranoside,<sup>12</sup>

sorbitan monolaurate exhibits interesting antimicrobial activity against Gram-positive bacteria. However, its activity could be seriously altered by the hydrolysis of the ester bond by cellular esterases, producing the corresponding fatty acid and the inactive polyol. For example, it has been shown that monolaurin can be quickly hydrolyzed ( $t_{1/2}$  of about 5 min) in the presence of *Staphylococcus aureus* cells.<sup>13</sup> In order to increase the life-time of antimicrobials, the ester linkage could be replaced by an ether group thus preventing such hydrolysis. For examples, 1-O-dodecylglycerol exhibits an enhanced activity against *Enterococcus faecium* than monolaurin<sup>14</sup> and methyl 6-O-dodecanyl- -D-glucopyranoside has an improved activity against *S. aureus* and *Listeria* spp. compared to its ester counterpart.<sup>12</sup> However, such approach has never been reported on sorbitan derivatives, probably due to the difficulty to prepare the corresponding ethers. In this context, we have previously prepared a range of sorbitan ethers<sup>15</sup> as robust analogues of span derivatives and we now report their antimicrobial activity against Gram-positive bacteria.<sup>16</sup>

A range of sorbitan ethers has been synthesized in two steps from sorbitan – a dehydration product of sorbitol – following an acetalization/hydrogenolysis sequence (Scheme 1).<sup>15</sup> First, acid-catalysed acetalization of non-protected sorbitan **1** with linear aldehydes gave a range of sorbitan acetals **2–6** with 29–81% isolated yields.<sup>17</sup> Noteworthy, these compounds exist as a mixture

\* Corresponding authors.

E-mail addresses: [gerard.lina@univ-lyon1.fr](mailto:gerard.lina@univ-lyon1.fr) (G. Lina), [nicolas.duguet@univ-lyon1.fr](mailto:nicolas.duguet@univ-lyon1.fr) (N. Duguet), [marc.lemaire.chimie@univ-lyon1.fr](mailto:marc.lemaire.chimie@univ-lyon1.fr) (M. Lemaire).



**Scheme 1.** Preparation of sorbitan ethers from sorbitan through acetalization/hydrogenolysis sequence.

**Table 1**

Antimicrobial activities of sorbitan acetals **2–6** and sorbitan ethers **7–11** against Gram positive bacteria.

Entry	Compounds	MIC values ( $\mu\text{g/mL}$ ) <sup>a</sup>		
		<i>L. monocytogenes</i> (CIP 103575)	<i>E. faecalis</i> (ATCC® 29212™)	<i>S. aureus</i> (ATCC® 29213™)
1	C5AcSorb <b>2</b>	> 512	> 512	> 512
2	C5EthSorb <b>7</b>	> 512	> 512	> 512
3	C6AcSorb <b>3</b>	> 512	> 512	> 512
4	C6EthSorb <b>8</b>	> 512	> 512	> 512
5	C8AcSorb <b>4</b>	> 512	> 512	> 512
6	C8EthSorb <b>9</b>	> 512	> 512	> 512
7	C10AcSorb <b>5</b>	> 512	> 512	> 512
8	C10EthSorb <b>10</b>	> 512	> 512	> 512
9	C12AcSorb <b>6</b>	8	8	32
10	C12EthSorb <b>11</b>	32	32	32
11	dodecanal	> 512	> 512	> 512

<sup>a</sup> MIC values were determined by serial dilution (2 by 2) of each compound from 512  $\mu\text{g/mL}$  to 0.25  $\mu\text{g/mL}$ .

of five-membered and six-membered isomers. Subsequent reductive cleavage of the mixture with hydrogen furnished the corresponding sorbitan ethers **7–11** with 55–85% isolated yields.<sup>17</sup> These species were obtained as an inseparable mixture of regioisomers with the alkyl chain grafted either at the position 3, 5 or 6. Noteworthy, this lack of selectivity is not detrimental for such low-added value compounds (Span 20 is actually a mixture of products).

The antimicrobial activity of sorbitan acetals (abbreviated CxAcSorb, with x = number of carbons of the alkyl chain, Ac = acetal, Sorb = sorbitan) and sorbitan ethers (abbreviated CxEthSorb, with x = number of carbons of the alkyl chain, Eth = ether, Sorb = sorbitan) were first evaluated against Gram-negative bacteria, namely, *Escherichia coli* (ATCC® 8739™) and *Pseudomonas aeruginosa* (ATCC® 27853™) by microdilution as previously described.<sup>18</sup> Unfortunately, no antimicrobial activity was observed for these bacteria (MIC >512  $\mu\text{g/mL}$ ). As a result, their antimicrobial activities were next evaluated against Gram-positive bacteria such as *Listeria monocytogenes* (CIP 103575), *Enterococcus faecalis* (ATCC® 29212™) and *S. aureus* (ATCC® 29213™). The results are gathered in Table 1.

No antimicrobial activity has been detected for sorbitan acetals **2–5** or ethers **7–10** bearing an alkyl chain with ten or less carbons (Table 1, entries 1–8). On the contrary, sorbitan acetal C12AcSorb **6** exhibits a significant antimicrobial activity against *S. aureus* (MIC = 32  $\mu\text{g/mL}$ ) and was found slightly more active against *L. monocytogenes* and *E. faecalis* (MIC = 8  $\mu\text{g/mL}$ ) (Table 1, entry 9). In that case, hydrolysis studies have been performed in order to determine whether the antimicrobial activity arises from the acetal derivative itself or from the hydrolyzed product. These studies were conducted at 40 °C at pH = 7 (physiological pH) but also at pH = 5 and 1 in order to get further information on the stability of such compounds (Fig. 1). At pH = 1, about 70% of the acetal was hydrolyzed after the first hour and the compound was totally degraded after 20 h. In sharp contrast, at pH = 7 or 5, only a slight degradation (about 4–6%) was observed after 1 h at 40 °C and about 92–94% of the compound still remains after 20 hours at this temperature. These results indicate that the rate of hydrolysis of acetal is slow under these conditions.

Moreover, the antimicrobial activity of dodecanal – the remaining amphiphilic product after hydrolysis – was also evaluated and no significant antimicrobial activity was observed (Table 1, entry

Download English Version:

<https://daneshyari.com/en/article/7780727>

Download Persian Version:

<https://daneshyari.com/article/7780727>

[Daneshyari.com](https://daneshyari.com)