

# Synthesis and biological evaluation of peptide-derived TSLP inhibitors



Seonghu Park, Yeeun Park, Sang-Hyun Son, Kiho Lee, Yong Woo Jung, Ki Yong Lee, Young Ho Jeon\*, Youngjoo Byun\*

College of Pharmacy, Korea University, 2511 Sejong-ro, Sejong 30019, South Korea

## ARTICLE INFO

### Article history:

Received 21 July 2017

Revised 31 August 2017

Accepted 1 September 2017

Available online 7 September 2017

### Keywords:

Allergic disease

TSLP inhibitors

Peptides

TSLP receptors

## ABSTRACT

Thymic stromal lymphopoietin (TSLP) is a type II cytokine which is associated with most inflammatory allergic disorders in humans. It is produced mainly by epithelial cells with important role in the development of chronic inflammatory diseases by activating T-helper cell type-2 ( $T_H2$ ) pathways. In this study, a total of 16 peptides were prepared by solid phase peptide synthesis based on amino acid sequences of the interface between TSLP and TSLP receptor. Their TSLP inhibition activities were determined by ELISA assay. Among them, three peptides (**6–8**) exhibited >50% inhibition at concentration of 0.3 mM. They can be used as hit compounds for developing peptide-based TSLP inhibitors.

© 2017 Elsevier Ltd. All rights reserved.

Allergic disease has increasingly emerged as a significant global health issue that affects millions of individuals worldwide.<sup>1</sup> Allergic disease is caused by unrestrained T helper 2 ( $T_H2$ )-biased immune responses that induce asthma, atopic dermatitis, and allergic rhinitis.<sup>2,3</sup> It has a major social and economic impact on patients, patients' family, and the society as a whole.<sup>4</sup> Despite being one of the most prevalent among all human disease categories, there is no effective vaccine or therapeutics to treat allergic diseases fundamentally. Therefore, understanding immune mechanisms involved in allergic diseases and finding treatment strategies for these diseases fundamentally are urgently needed.

Thymic stromal lymphopoietin (TSLP) is a cytokine belonging to interleukin 2 (IL-2) family. It is expressed mostly by epithelial cells on barrier surfaces such as the skin, lung, and gut in response to external stimuli.<sup>5–7</sup> TSLP is a key initiator of STAT5-mediated  $T_H2$  inflammatory pathways.<sup>3,8–10</sup> It exerts biological functions by making triplex structure with an interleukin-7 (IL-7) receptor  $\alpha$  chain (IL-7R $\alpha$ ) and a unique TSLP receptor (TSLPR).<sup>11,12</sup> TSLP alone binds to IL-7R $\alpha$  with comparatively low affinity ( $K_d = 2.3 \mu\text{M}$  in mouse and no affinity was detected at 100 nM of IL-7R $\alpha$  in human). TSLP-mediated  $T_H2$  signaling is explained by sequential and cooperative formation of the ternary complex (TSLP/TSLPR/IL-7R $\alpha$ ). The TSLP-TSLPR binary complex ( $K_d = 58 \text{ nM}$  in mouse and  $K_d = 32 \text{ nM}$  in human) is formed first and then establishes the stable ternary complex with IL-7R $\alpha$ .<sup>13</sup> Heterodimerization of TSLP with TSLPR enhanced its binding affinity for IL-7R $\alpha$  ( $K_d = 1.5 \text{ nM}$  in mouse


and 29 nM in human).<sup>13,14</sup> Forming a triple complex of TSLP/TSLPR/IL-7R $\alpha$  activates JAK1 and JAK2, thereby resulting in phosphorylation of STAT5.<sup>15</sup>

Numerous reports have highlighted the fact that aberrant TSLP signaling is closely associated with inflammatory allergic diseases, including asthma, atopic dermatitis, chronic obstructive pulmonary disease (COPD), and eosinophilic esophagitis.<sup>8,9,16–18</sup> For instance, high TSLP expression levels are known to be correlated with the severity of allergic diseases in human and mice.<sup>3,9,19</sup> Another study has shown that blocking TSLPR in a primate animal model can result in resistance to the development of allergic inflammation.<sup>20</sup> Recently, strategy for targeting TSLP, IL-25, and IL-33 together has demonstrated therapeutic potential in mouse disease models of inflammation and fibrosis.<sup>21</sup> Moreover, blocking TSLP signaling with an anti-TSLP monoclonal antibody is under investigation in a clinical trial to treat asthma patients.<sup>22</sup> These results suggest that TSLP represents a key mediator in the pathogenesis of allergic disease. Therefore, blocking TSLP signaling is considered as an attractive intervention strategy to allergic diseases fundamentally.

Identifying potent and selective inhibitors of TSLP as pharmacological tools to better understand their roles in allergic inflammatory response and their potential as therapeutics of allergic disease has been one of our research interests. Recently, Savvides and co-workers have reported the first high-resolution X-ray structure of mouse and, more recently, human TSLP/TSLPR/IL-7R $\alpha$  complexes.<sup>13,14</sup> The 3D X-ray crystal structure of TSLP/TSLPR/IL-7R $\alpha$  complex provides insight into its structure-biological function relationship. It assists in the design of TSLP inhibitors for drug

\* Corresponding authors.

E-mail addresses: [yhjeon@korea.ac.kr](mailto:yhjeon@korea.ac.kr) (Y.H. Jeon), [yjbyun1@korea.ac.kr](mailto:yjbyun1@korea.ac.kr) (Y. Byun).



**Fig. 1.** Design of peptide-derived hTSLP inhibitors based on amino acid sequences of mTSLPR in the interface site between mTSLP and mTSLPR. SET 1 and SET 2 consist of ARQRASAW and RAAHYGLEA sequences, respectively.


development to treat allergic diseases. Since there has been no report on TSLP inhibitors based on peptides or small molecules, the objective of this study was to discover novel peptide-based TSLP inhibitors, by utilizing structural information of the interface between TSLP and TSLPR.

Previously, it has been reported that mouse TSLP (mTSLP) can interact with mouse TSLPR (mTSLPR) and mouse IL-7R $\alpha$  heterodimer complex via two equally extensive interaction interfaces.<sup>13</sup> Furthermore, binding of mTSLP to mTSLPR has been characterized by the formation of two specific interaction sites: amino acid residues of 107–114 (SET 1) and 191–199 (SET 2) of mTSLPR (Fig. 1). Based on these interactions between mTSLP and mTSLPR, our hypothesis was that mimicking these amino acid residues of these specific interaction sites could disrupt the interaction between TSLP and TSLPR. As an initial step of peptide-based inhibitor design, we have exploited the mTSLPR sequence because there was the only mTSLP–mTSLPR structure available in Protein Data Bank (PDB ID: 4NN5) when we designed this experiment.<sup>13</sup> The sequence identity between hTSLPR and mTSLPR was 36.1%, and the similarity was 49.2%. The corresponding sequences of SET1

and SET2 in human TSLPR are ASRWNVYY and MEDVYGPDT, respectively.

Peptide-based inhibitors of protein-protein interactions (PPI) have several advantages, including high structural similarity to fragments of target protein, facile synthesis by using solid-phase peptide synthesis (SPPS), and opportunity to modify peptide sequences with various functional groups.<sup>23</sup> Even small linear peptides by mimicking epitopes of TSLP–TSLPR interaction sites can serve as a starting point for the design of TSLP inhibitors. Herein, we first report the synthesis of peptide-based TSLP inhibitors using SPPS and evaluation of their biological activities using ELISA assay.

To verify our hypothesis, peptides (**1–8**) derived from SET 1 and peptides (**9–16**) from SET 2 were synthesized. SPPS was applied in the synthesis of target peptides by using Fmoc-protected amino acid resin as starting materials (Scheme 1). Peptide coupling reactions were performed by using HBTU (3.0 eq)/HOBt (2.0 eq)/DIPEA (6.0 eq) in DMF for 1 h. Fmoc group in each step was removed by applying 20% piperidine in DMF for 20 min. Acetylation of *N*-terminal amine group was accomplished by treating acetic anhydride and DIPEA in DMF at room temperature. In the final step of



**Scheme 1.** Synthesis of peptide-derived TSLP inhibitors through solid-phase peptide synthesis (SPPS). Reagents and condition: (a) (i) 20% piperidine/DMF, 20 min; (ii) Fmoc-Aa2-OH, HBTU, DIPEA, HOBt, DMF, 1 h; (b) (i) 20% piperidine/DMF, 20 min, rt; (ii) Fmoc-Aa3-OH, HBTU, DIPEA, HOBt, DMF, 1 h; (c) 20% piperidine/DMF, 20 min; (d) TFA/thioanisole/H<sub>2</sub>O/TIS (95:2:2:1, v/v/v/v), 3 h; (e) Ac<sub>2</sub>O, DIPEA, DMF, 1 h.

Download English Version:

<https://daneshyari.com/en/article/7780787>

Download Persian Version:

<https://daneshyari.com/article/7780787>

[Daneshyari.com](https://daneshyari.com)