

Accepted Manuscript

Title: Two-step sequential cross-linking of sugar beet pectin for transforming zein nanoparticle-based Pickering emulsions to emulgels

Author: Sahar Soltani Ashkan Madadlou



PII: S0144-8617(15)00965-0
DOI: <http://dx.doi.org/doi:10.1016/j.carbpol.2015.09.100>
Reference: CARP 10402

To appear in:

Received date: 24-6-2015
Revised date: 31-8-2015
Accepted date: 26-9-2015

Please cite this article as: Soltani, S., and Madadlou, A., Two-step sequential cross-linking of sugar beet pectin for transforming zein nanoparticle-based Pickering emulsions to emulgels, *Carbohydrate Polymers* (2015), <http://dx.doi.org/10.1016/j.carbpol.2015.09.100>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Research highlights

- Zein particles with mean size of 64 nm were formed by solvent evaporation.
- Several zein particles-based and pectin-enriched Pickering emulsions were prepared.
- The emulsions were charged sequentially with laccase and CaCl_2 yielding emulgels.
- The emulgels were self-standing and became harder with increasing pectin content.

Download English Version:

<https://daneshyari.com/en/article/7786723>

Download Persian Version:

<https://daneshyari.com/article/7786723>

[Daneshyari.com](https://daneshyari.com)