


Accepted Manuscript

Title: Distribution of phosphorus and hydroxypropyl groups within granules of modified sweet potato starches as determined after chemical peeling

Author: Jianwei Zhao Henk A Schols Zhenghong Chen
Zhengyu Jin Piet Buwalda Harry Gruppen



PII: S0144-8617(15)00587-1
DOI: <http://dx.doi.org/doi:10.1016/j.carbpol.2015.06.078>
Reference: CARP 10069

To appear in:

Received date: 22-2-2015
Revised date: 7-6-2015
Accepted date: 23-6-2015

Please cite this article as: Zhao, Jianwei., Schols, Henk A., Chen, Zhenghong., Jin, Zhengyu., Buwalda, Piet., & Gruppen, Harry., Distribution of phosphorus and hydroxypropyl groups within granules of modified sweet potato starches as determined after chemical peeling. *Carbohydrate Polymers* <http://dx.doi.org/10.1016/j.carbpol.2015.06.078>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Distribution of phosphorus and hydroxypropyl groups within granules of modified sweet potato starches as determined after chemical peeling

Jianwei Zhao^{a,b}, Henk A Schols^a, Zhenghong Chen^c, Zhengyu Jin^b, Piet Buwalda^c, Harry Gruppen^{a,*}

^a Wageningen University, Laboratory of Food Chemistry, P.O. Box 17, 6700AA Wageningen, The Netherlands

^b Jiangnan University, School of Food Science and Technology, 214122 Wuxi, P.R. China

^c AVEBE Food Innovation Centre, Veendam, The Netherlands

*Corresponding author

Prof., Dr. Harry Gruppen

Wageningen University, Laboratory of Food Chemistry, P.O. Box 17,
6700AA Wageningen, The Netherlands

Tel: +31 317 483211

E-mail: Harry.Gruppen@wur.nl

Highlights

- The starch granule was peeled gradually by calcium chloride and enzymes.
- Cross-linking and hydroxypropylation occur predominantly in the granule periphery.
- Phosphorus levels in different parts of the large-size granules varied.

Download English Version:

<https://daneshyari.com/en/article/7787643>

Download Persian Version:

<https://daneshyari.com/article/7787643>

[Daneshyari.com](https://daneshyari.com)