

Boronate affinity monolith via two-step atom transfer radical polymerization for specific capture of *cis*-diol-containing compounds

Qian Dong, Shuai-Shuai Chi, Xi-Yan Deng, Yao-Han Lan, Chao Peng, Lin-Yi Dong, Xian-Hua Wang

PII: S0014-3057(17)31703-2
DOI: https://doi.org/10.1016/j.eurpolymj.2018.02.007
Reference: EPJ 8281

To appear in: *European Polymer Journal*

Received Date: 24 September 2017
Revised Date: 5 February 2018
Accepted Date: 5 February 2018

Please cite this article as: Dong, Q., Chi, S-S., Deng, X-Y., Lan, Y-H., Peng, C., Dong, L-Y., Wang, X-H., Boronate affinity monolith via two-step atom transfer radical polymerization for specific capture of *cis*-diol-containing compounds, *European Polymer Journal* (2018), doi: https://doi.org/10.1016/j.eurpolymj.2018.02.007

Boronate affinity monolith via two-step atom transfer radical polymerization for specific capture of *cis*-diol-containing compounds

Qian Dong[1], Shuai-Shuai Chi[1], Xi-Yan Deng, Yao-Han Lan, Chao Peng, Lin-Yi Dong, Xian-Hua Wang*

Tianjin Key Laboratory on Technologies Enabling Development of Clinical Therapeutics and Diagnostics, School of Pharmacy, Tianjin Medical University, Tianjin 300070, P. R. China.

[1] These authors contributed equally to this work.

*** Corresponding author:** Dr. Xian-Hua Wang,

Postal address: Building B for School of Pharmacy, Tianjin Medical University, 22 Qixiangtai Road, Heping District, Tianjin 300072, China

Tel.: +86-022-83336675; Fax.: +86-022-83336690;

E-mail: xianhua.w@163.com

Keywords: Boronate affinity; *cis*-Diol-containing compounds; Porous monolith; Atom-transfer radical polymerization;