

Contents

LETTERS

NMR-based metabolomics study of *Amphoricarpus* species from Montenegro

Phytochemistry Letters 25 (2018) pp. 1–5


Mirjana Cvetković^a, Boban Andelković^b, Vladimir Stevanović^c, Milka Jadranin^a, Iris Đorđević^{d,*}, Vele Tešević^b, Slobodan Milosavljević^c, Dejan Godevac^a

^aInstitute of Chemistry, Technology and Metallurgy, University of Belgrade, Studentski trg 12-16, 11000 Belgrade, Serbia

^bFaculty of Chemistry, University of Belgrade, Studentski trg 12-16, 11000 Belgrade, Serbia

^cSerbian Academy of Sciences and Arts, Knez Mihailova 35, 11000 Belgrade, Serbia

^dFaculty of Veterinary Medicine, University of Belgrade, Bulevar oslobođenja 18, 11000 Belgrade, Serbia

Bioactive triterpenoid glycosides from the twigs and leaves of *Camellia reticulata*

Phytochemistry Letters 25 (2018) pp. 6–11

Qi-Run Chen^{a,c,1}, Dan Li^{a,d,1}, Xu-Jie Qin^a, Hui Liu^a, Wei Ni^a, Huan Yan^a, Li He^{b,*}, Qing Zhao^c, Xiao Ma^c, Hai-Yang Liu^{a,*}


^aState Key Laboratory of Phytochemistry and Plant Resources in West China, and Yunnan Key Laboratory of Natural Medicinal Chemistry, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, China

^bDepartment of Dermatology, The First Affiliated Hospital of Kunming Medical University, Kunming 650032, China

^cYunnan University of Traditional Chinese Medicine, Kunming 650500, China

^dUniversity of Chinese Academy of Sciences, Beijing 100039, China

^eKunming Botanical Bio-Technique Co. Ltd., Kunming 650106, China

Garcixanthones B and C, new xanthones from the pericarps of *Garcinia mangostana* and their cytotoxic activity

Phytochemistry Letters 25 (2018) pp. 12–16

Sabrin R.M. Ibrahim^{a,b,*}, Hossam M. Abdallah^{c,d}, Ali M. El-Halawany^d, Mohammed F. Radwan^{e,f}, Ibrahim A. Shehata^{c,d}, Essam M. Al-Harshany^c, Mohamed F. Zayed^a, Gamal A. Mohamed^{c,g}

^aDepartment of Pharmacognosy and Pharmaceutical Chemistry, College of Pharmacy, Taibah University, Al Madinah Al Munawwarah, 30078, Saudi Arabia

^bDepartment of Pharmacognosy, Faculty of Pharmacy, Assiut University, Assiut, 71526, Egypt


^cDepartment of Natural Products and Alternative Medicine, Faculty of Pharmacy, King Abdulaziz University, Jeddah, 21589, Saudi Arabia

^dDepartment of Pharmacognosy, Faculty of Pharmacy, Cairo University, Cairo, 11562, Egypt

^eDepartment of Pharmaceutical Chemistry, Faculty of Pharmacy, King Abdulaziz University, Jeddah, 21589, Saudi Arabia

^fDepartment of Medicinal Chemistry, Faculty of Pharmacy, Minia University, Minia, 61519, Egypt

^gPharmacognosy Department, Faculty of Pharmacy, Al-Azhar University, Assiut Branch, Assiut, 71524, Egypt



Acylated iridoid diglycosides from the cultivated endangered ornamental tree *Gmelina hainanensis*

Phytochemistry Letters 25 (2018) pp. 17–21


Juan Xiong^a, Xi-Ying Wu^a, Pei-Pei Wang^b, Chaiwan Lau^{c,*}, Hui Fan^d,
Guang-Lei Ma^a, Yu Tang^a, Jia Li^b, Jin-Feng Hu^{a,*}

^aDepartment of Natural Products Chemistry, School of Pharmacy, Fudan University, Shanghai, 201203, PR China

^bState Key Laboratory of Drug Research, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, Shanghai, 201203, PR China

^cInstrumental Analysis Center, School of Pharmacy, Fudan University, Shanghai, 201203, PR China

^dShanghai Key Laboratory of Green Chemistry and Chemical Processes, School of Chemistry and Molecular Engineering, East China Normal University, Shanghai, 200062, PR China



Pahangine A and B, two new oxetane containing neolignans from the barks of *Beilschmiedia glabra* Kosterm (Lauraceae)

Phytochemistry Letters 25 (2018) pp. 22–26

Syazreen Nadia Sulaiman^a, Azeana Zahari^{a,b}, Sook Yee Liew^c, Marc Litaudon^d, A.M. Issam^a,
Habibah A. Wahab^e, Khalijah Awang^{a,b,*}


^aDepartment of Chemistry, Faculty of Science, University of Malaya, 50603 Kuala Lumpur, Malaysia

^bCentre of Natural Products and Drug Discovery (CENAR), University of Malaya, 50603 Kuala Lumpur, Malaysia

^cChemistry Division, Centre for Foundation Studies in Science, University of Malaya, 50603 Kuala Lumpur, Malaysia

^dCentre de Recherche de Gif, Institut de Chimie des Substances Naturelles, CNRS-ICSN UPR 2361, Université Paris-Saclay, Avenue de la Terrasse, 91198 Gif-sur-Yvette Cedex, France

^eSchool of Pharmaceutical Sciences, Universiti Sains Malaysia, 11800 Universiti Sains Malaysia, Pulau Pinang, Malaysia




Monoterpene, benzyl and 3,4-dihydroxyphenethyl glycosides from *Magnolia thailandica*

Phytochemistry Letters 25 (2018) pp. 28–32

Tripetch Kanchanapoom^{a,b,*}, Poolsak Sahakitpichan^a, Nitirat Chimnoi^a,
Chutima Srinroch^a, Wassapol Thamniyom^a, Somsak Ruchirawat^a

^aChulabhorn Research Institute, Kamphaeng Phet 6, Talat Bang Khen, Lak Si, Bangkok, 10210, Thailand

^bFaculty of Pharmaceutical Sciences, Khon Kaen University, Khon Kaen, 40002, Thailand



Protective effects of dammarane-type triterpenes from hydrolyzate of *Gynostemma pentaphyllum* against H₂O₂-induced injury and anti-hepatic fibrosis activities


Phytochemistry Letters 25 (2018) pp. 33–36

Xiaoshu Zhang^{a,b}, Guohui Shi^{a,b}, Mingbing Liu^a, Rong Chen^a, Xiaojun Wu^c, Yuqing Zhao^{a,b,*}

^aSchool of Functional Food and Wine, Shenyang Pharmaceutical University, Shenyang, 110016, People's Republic of China

^bKey Laboratory of Structure-Based Drug Design & Discovery, Ministry of Education, Shenyang Pharmaceutical University, Shenyang, 110016, People's Republic of China

^cShenyang Pharmaceutical University, Shenyang, 110016, People's Republic of China



Download English Version:

<https://daneshyari.com/en/article/7818265>

Download Persian Version:

<https://daneshyari.com/article/7818265>

[Daneshyari.com](https://daneshyari.com)