



First total synthesis of the anti-inflammatory and pro-resolving lipid mediator 16(*R*),17(*S*)-diHDHA

Ana R. Rodriguez, Bernd W. Spur^{*}

Department of Cell Biology and Neuroscience, Rowan University-SOM, Stratford, NJ 08084, USA

ARTICLE INFO

Article history:

Received 22 January 2018

Revised 3 February 2018

Accepted 12 February 2018

Available online 13 February 2018

Dedicated to Professor E. J. Corey on the occasion of his 90th birthday.

Keywords:

16(*R*),17(*S*)-diHDHA

Total synthesis

Inflammation resolution

Base-promoted epoxide isomerization

Lipoxidase

Chiral pool

ABSTRACT

The first total synthesis of the anti-inflammatory and pro-resolving lipid mediator 16(*R*),17(*S*)-diHDHA, derived from docosahexaenoic acid (DHA), and its 16-epimer have been achieved. Two synthetic approaches are described for the synthesis of 16(*R*),17(*S*)-diHDHA. The first strategy started from DHA and used an enzymatic reaction, a vanadium catalyzed allylic epoxidation and a base-promoted epoxide isomerization. The second approach utilized a chiral pool strategy starting from 2-deoxy-D-ribose to establish the chiral centers; Wittig reactions, mild acetonide cleavage and ester hydrolysis were the key steps in the synthesis.

© 2018 Elsevier Ltd. All rights reserved.

Inflammation is a response to tissue damage and/ or infection. Acute inflammation that is self-limited and achieves resolution is a protective mechanism,^{1,2} but uncontrolled inflammation is the underlining course of chronic diseases including autoimmune diseases, cancer, diabetes, asthma and neurological diseases such as Alzheimer's and Parkinson disease.³ It is well documented that ω-3 fatty acids mainly found in fish oil have anti-inflammatory properties and help with the resolution of inflammation.^{4–6} The pioneering work by Serhan and collaborators revealed that these ω-3 poly-unsaturated fatty acids are converted at the site of inflammation by lipoxygenase enzymes to potent lipid mediators that are responsible for the resolution of inflammation.^{7,8}

Several families of pro-resolving lipid mediators have been identified and collectively named specialized pro-resolving mediators (SPMs): lipoxins derived from arachidonic acid, resolvins of the E-type derived from eicosapentaenoic acid (EPA), resolvins of the D-type, maresins, protectins and their sulfido-conjugates derived from docosahexaenoic acid (DHA) and the more recently uncovered metabolites from n-3 docosapentaenoic acid (n-3 DPA).^{9–16}

The proposed biosynthesis of the protectins is outlined in Fig. 1.¹⁷ DHA is converted by the 15-lipoxidase to 17(*S*)-hydroperoxy-docosahexaenoic acid [17(*S*)-HpDHA] that undergoes enzymatic conversion to the allylic epoxide 16(*S*),17(*S*)-epoxy-docosahexaenoic acid. This intermediate undergoes enzymatic hydrolysis to 10(*R*),17(*S*)-protectin D1 (PD1) whereas a soluble epoxide hydrolase gives the 16(*R*),17(*S*)-diHDHA. The same 16(*S*),17(*S*)-epoxy-docosahexaenoic acid intermediate can be converted by a glutathione S-transferase via a SN2 mechanism to the Protectin conjugate in tissue regeneration (PCTR).^{18–20} Due to the growing interest to study the biological and pharmacological properties of these mediators combined with the limited availability from natural sources these products have to be prepared by chemical synthesis. We have recently reported the total synthesis of PD1²¹ and the PCTRs.¹⁹ In the current publication we wish to report the total synthesis of 16(*R*),17(*S*)-diHDHA [(4*Z*,7*Z*,10*Z*,12*E*,14*E*,16*R*,17*S*,19*Z*)-16,17-dihydroxy-4,7,10,12,14,19-docosahexaenoic acid (**1**)].

As shown in the retrosynthetic scheme (Fig. 2) we have prepared 16(*R*),17(*S*)-diHDHA via two different strategies. In the first approach we used an enzymatic-chemical route starting from DHA (**3**). In the second approach we employed a chiral pool strategy starting from 2-deoxy-D-ribose. The key intermediates **4** and **5** were used to produce the 16(*R*),17(*S*)-diHDHA skeleton.

^{*} Corresponding author.

E-mail address: spurbw@rowan.edu (B.W. Spur).




Fig. 1. Proposed biosynthesis of 10R,17S-Protectin D1, 16R,17S-diHDHA and PCTRs.




Fig. 2. Two retrosynthetic approaches to 16(R),17(S)-diHDHA.

As shown in Scheme 1 DHA (3) was reacted with lipoxidase from soybean in borate buffer pH 9 in the presence of O₂ followed by reduction with Ph₃P to obtain 17(S)-hydroxy-docosahexaenoic acid [17(S)-HDHA] (2) with >97.2% ee as determined by chiral-HPLC (racemic 17-HDHA was prepared from 2 by Dess-Martin oxidation followed by NaBH₄ reduction in ethanol).²² Esterification of 2 with trimethylsilyldiazomethane in CH₃OH/toluene afforded compound 11 in 85% isolated yield²² that was submitted to allylic epoxidation with TBHP in the presence of a catalytic amount of vanadyl acetylacetonate to afford a mixture of *erythro* and *threo* epoxy-alcohols 12 and 13 in the proportion of 77/23 respectively.²³ It should be mentioned that excess TBHP had to be avoided to min-

imize diepoxidation. The separation of the epoxides 12 and 13 was easily achieved by flash chromatography in the presence of 1% Et₃N providing a convenient access to 16(R),17(S)-diHDHA (1) and its 16-*epi*-isomer 16 respectively. Compounds 12 (*erythro*) and 13 (*threo*) had the characteristic ¹H–¹H coupling constants between the epoxide and the alcohol (*J*_{16,17} = 3.0 Hz for 12 and 4.5 Hz for 13).^{23,24} Mild hydrolysis of the methyl ester in compound 12 with 1 N LiOH in THF/CH₃OH/H₂O 4/1/1 followed by acidification with sat. NaH₂PO₄ in the presence of ethyl acetate afforded *erythro* 15 (S),16(S)-epoxy-17(S)-HDHA (14) that was submitted to a base-promoted epoxide isomerization, modified from the procedures reported by the groups of Falck and Corey.^{25–28} The best results were obtained by adding 5 equiv of KHMDS in toluene to 14 in THF at 0 °C affording 16(R),17(S)-diHDHA (1). The geometry of the 10Z,12E,14E-triene in 1 (*J*_{10,11} = 11.1, *J*_{12,13} = 14.4 Hz and *J*_{14,15} = 15.0 Hz) was confirmed by the ¹H–¹H coupling constants.²⁴ The ¹H NMR, ¹³C NMR, UV and HPLC/UV/MS analysis were consistent with the structures of 1.²⁴ In a similar manner *threo*-13 was converted to 16(S),17(S)-diHDHA (16).²⁴

In the second strategy the key intermediate 4 was prepared from compound 17 (Scheme 2). We had previously reported the synthesis of 17 from the intermediates 6 and 7.¹⁹ The free hydroxy group in compound 17 was converted to its tosylate with 3 equiv of tosyl chloride in CH₂Cl₂ in the presence of pyridine to afford 18, which was then transformed to the phosphonium salt 4 with triphenylphosphine in acetonitrile at reflux in 84% yield.

The C11–C22 chiral fragment 5 was obtained in 4 steps from 3,4-*O*-isopropylidene-2-deoxy-d-ribose (9) (Scheme 3).²⁹ Wittig reaction of 9 with 2.5 equiv of the ylide generated in situ from *n*-propyltriphenylphosphonium bromide (10) with *n*-BuLi (2.5 equiv) in THF in the presence of HMPA at –78 °C produced 19 in 76% yield.³⁰ Compound 19 was converted to the aldehyde 20 by oxidation with PCC in CH₂Cl₂.³¹ Four-carbon homologation of the aldehyde 20 with recrystallized (2E)-4-triphenylphosphoranylidene-2-butenal (8)^{32,33} in CH₂Cl₂ followed by treatment with catalytic iodine in benzene afforded the *E,E*-dienal 5 in 35% yield (2-steps).

Wittig reaction of the phosphorane prepared in situ from the phosphonium tosylate 4 with KHMDS at –78 °C in THF with the dienal 5 gave the isopropylidene protected 16(R),17(S)-diHDHA methyl ester (21) in 52% yield based on recovered aldehyde 5 (Scheme 3). The geometry of the 12E,14E-diene unit in 5 (*J*_{12,13} = 15.3 Hz and *J*_{14,15} = 15.3 Hz) and the 10Z,12E,14E-triene in 21 (*J*_{10,11} = 10.8 Hz, *J*_{12,13} = 14.4 Hz and *J*_{14,15} = 15.0 Hz) were confirmed by the ¹H–¹H coupling constants.²⁴ The isopropylidene protective group was cleaved with *p*-toluenesulfonic acid in CH₃OH at 0 °C to give 16(R),17(S)-diHDHA methyl ester (22). Mild hydrolysis of 22 with 1 N LiOH in CH₃OH/H₂O at 0 °C under argon followed by acidification with sat. NaH₂PO₄ (pH 3) in the presence of ethyl acetate, afforded 16(R),17(S)-diHDHA (1) in 96% yield. Co-injection of 16(R),17(S)-diHDHA (1) prepared from both routes were analyzed by HPLC/UV/MS and found to be identical.

In summary we have developed two routes for the synthesis of 16(R),17(S)-diHDHA (1),²⁴ making this anti-inflammatory and pro-resolving lipid mediator from docosahexaenoic acid available for further biological and pharmacological testing. The synthesis of the epimeric 16(S),17(S)-diHDHA (16)²⁴ has also been accomplished. The synthesis of other specialized pro-resolving mediators (SPMs) will be reported in due course.

Acknowledgments

Financial support of this research in part by the NJHF and NJCBIR is gratefully acknowledged.

Download English Version:

<https://daneshyari.com/en/article/7830121>

Download Persian Version:

<https://daneshyari.com/article/7830121>

[Daneshyari.com](https://daneshyari.com)