



Daucus carota root enzyme catalyzed Henry reaction: A green approach

Chiranjit Acharya, Anushree Achari, Parasuraman Jaisankar*

Laboratory of Catalysis and Chemical Biology, Department of Organic and Medicinal Chemistry, CSIR-Indian Institute of Chemical Biology, 4 Raja S. C. Mullick Road, Jadavpur, Kolkata 700 032, India

ARTICLE INFO

Article history:

Received 4 December 2017

Revised 30 December 2017

Accepted 4 January 2018

Available online 5 January 2018

Keywords:

Daucus carota

Enzyme

Henry reaction

Nitromethane

β -Nitroalcohol

ABSTRACT

Enzyme from *Daucus carota* root catalyzed Henry reaction of substituted benzaldehydes and nitromethane in phosphate buffer of pH 7 at 28 °C to afford β -nitroalcohols in excellent yields (up to 94%).

© 2018 Elsevier Ltd. All rights reserved.

The hidden art of organic synthesis is the carbon–carbon (C–C) bond forming reactions, the fundamental tool for the synthesis of complex molecular architecture.^{1–4} A number of methodologies have been reported over the years for C–C bond formation.^{5–9} Among various other protocols aldol reactions are the most common and well established methods.¹⁰ Besides this, Henry reaction of aldehydes with nitroalkanes to afford β -nitroalcohols which had been reported in 1895, has attracted the attention of medicinal chemists due to its applications in the synthesis of pharmaceutical agents,¹¹ precursors of natural products¹¹ and bioactive compounds having medicinal importance (Fig 1).¹²

The use of base in Henry reactions resulted in side products along with the desired β -nitroalcohol.¹³ Although, this shortcoming has been addressed by the use of metal and organo catalysts in place of conventional bases,¹⁴ the problem of toxicity and high cost is a deterrent. Therefore, further development was needed to provide green and efficient methodologies.

As a result of this development, enzyme promiscuity has been introduced.¹⁵ But only a few enzymes which are both costly and substrate specific, were reported to catalyze Henry reactions.^{16,17} Hence, the search for new enzymes provoked the development of enzyme catalysis which is an emerging field of modern organic synthesis.

Previously, Griengl et al. reported hydroxynitrile lyases (HNLs, protease enzyme) catalyzed synthesis of β -hydroxynitriles and β -nitroalcohols from nitrobenzaldehydes and hydrogencyanide

(HCN) or nitromethane with high substrate specificity.¹⁵ The primary role of HNLs is to abstract the acidic proton from HCN ($pK_a = 9.2$) and nitromethane ($pK_a = 10.21$). There were few reports available in the literature on *Daucus carota* root mediated reduction of optically active alcohols,¹⁸ until our recent report on its use as catalyst for the asymmetric cross aldol reaction of nitrobenzaldehyde and acetone.¹⁹ In continuation of our prior work¹⁹ we have deliberately used *Daucus carota* root or its enzyme in the promiscuous mixture of substituted nitrobenzaldehyde and nitromethane in aqueous medium to afford β -nitroalcohol in excellent yields (Scheme 1).

Initially the reaction was performed by stirring 2-nitrobenzaldehyde (**1a**, 1 mmol) and nitromethane (**2**, 1 mmol) in a suspension of freshly cut *Daucus carota* root (5.0 g) in aqueous medium for 24 h, which resulted in the formation of β -nitroalcohol (**3a**) with 55% of isolated yield (Table 1, entry 2). Encouraged by this initial result we then performed the same reaction with purified enzyme (10 μ L) from *Daucus carota* root (see SI) and it afforded the desired β -nitroalcohol with 70% of isolated yield in 15 h. Subsequently we went for optimizing the reaction protocol by changing reaction parameters (Table 1). No reaction took place in the absence of enzyme.

The optimized condition of the *Daucus carota* root enzyme catalyzed Henry reaction of 2-nitrobenzaldehyde and nitromethane to afford β -nitroalcohol in excellent yield (93%) was found to be phosphate buffer of pH 7 at 28 °C in 8 h (Table 1, entry 5). Organic solvents did not have any positive influence in this reaction, possibly, due to denaturation of the enzyme occurred in organic solvents. A number of β -nitroalcohols have been synthesized under the

* Corresponding author.

E-mail address: jaisankar@iicb.res.in (P. Jaisankar).





Fig. 1. β -Aminoalcohols having profound medicinal importance.



Scheme 1. *Daucus carota* root catalyzed Henry reaction.


optimized conditions and the results are summarized in Table 2. All the synthesized β -nitroalcohols, which were racemic in nature have been characterized by ^1H NMR, ^{13}C NMR, ESI-HRMS and FT-IR spectroscopic techniques (see SI).

It was observed that there was almost no electronic (viz. entry 3 vs. 6) influence of aryl substituents on the yields of the products. As one would expect, the high specificity of any enzyme in catalytic activity was corroborated when nitroethane and nitropropane were used in order to broaden the scope of the reaction, failed to undergo this reaction. We also checked the reactivity of aliphatic aldehydes with nitromethane in this promiscuous reaction but disappointingly, no reaction took place. The reusability of the *Daucus carota* root enzyme was investigated in the reaction of **1a** and **2** to yield **3a** which was extracted in diethylether followed by subsequent addition of the substrates (**1a** and **2**) in the aqueous layer which contains enzyme to catalyze next batches. It was perceived that the enzyme could catalyze the Henry reaction of **1a** and **2** up to five cycles to afford **3a**, though with progressively lower isolated yields (Table 3).

Earlier, it was reported that the lysine residue of HNLs acted as base to abstract a methylene proton from nitromethane to favor the reaction with nitrobenzaldehyde.^{15b} Recently, we disclosed that the *Daucus carota* root enzyme also contains Lysine residue which was proposed to favor the asymmetric cross aldol reaction of nitrobenzaldehyde and acetone.¹⁹ Therefore, we could visualize that the catalytic site of the enzyme¹⁹ which may contain lysine residue could activate the methylene proton of nitromethane (**2**) to favor Henry reaction with 2-nitrobenzaldehyde (**1a**), that resulted in the formation of β -nitroalcohol (**3a**) (Scheme 2). The non-stereospecificity of the products (**3a–i**) could be due to lack of binding of the substrate (**2**) with the amino acid residues present in the active sites of enzyme.

In conclusion we have disclosed yet another catalytic activity of *Daucus carota* root enzyme towards Henry reaction in aqueous medium. The impact of the reaction conditions including solvent, pH of the reaction medium has been explored and extent of the reaction investigated. The substrate specificity of the enzyme may be exploited in the synthesis of β -nitroalcohols which are important intermediates of many pharmaceuticals and natural products. The present methodology has some genuine and competitive advantages over the reported ones, including simple and mild

Table 1
Optimization of parameters for the *Daucus carota* root enzyme catalyzed Henry reaction^a



Entry	Solvent	Temperature ⁱ (°C)	Time ^γ (h)	Isolated yield (%)
1 [#]	Water	28	24	No reaction
2 ^φ	Water	28	24	55
3	Water	28	15	70
4	Phosphate buffer of pH 8	28	8	90
5 [§]	Phosphate buffer of pH 7	28	8	93
6	Phosphate buffer of pH 6	28	8	82
7	Phosphate buffer of pH 5	28	8	60
8	Diethylether (DEE)	28	8	No reaction
9	DEE: buffer of pH 7 (1:1)	28	12	50
10	TBME ^ε	28	8	No reaction
11	TBME: buffer of pH 7 (1:1)	28	12	60
12	Acetonitrile: buffer of pH 7 (1:1)	28	24	No reaction
13	Ethanol: buffer of pH 7 (1:1)	28	24	No reaction
14	Methanol: buffer of pH 7 (1:1)	28	24	No reaction

^a See experimental section for general reaction procedure.

[#] Reaction was placed without *Daucus carota* root enzyme.

^φ Reaction was placed with homogenized carrot root (5 g).

ⁱ The reaction was performed at 28 °C as the enzyme was reported to perform best at that temperature.¹⁹

^γ Further increase of reaction time has no effect on the yield of the product.

^ε *t*-Butylmethylether.

[§] Optimized condition.

Download English Version:

<https://daneshyari.com/en/article/7830609>

Download Persian Version:

<https://daneshyari.com/article/7830609>

[Daneshyari.com](https://daneshyari.com)