

Accepted Manuscript

Density functional reactivity theory characterizing the reactivity of frustrated Lewis pairs

Dongling Wu, Anjie Liu, Dianzeng Jia

PII: S2210-271X(18)30109-9
DOI: <https://doi.org/10.1016/j.comptc.2018.03.027>
Reference: COMPTC 2761

To appear in: *Computational & Theoretical Chemistry*

Received Date: 9 January 2018
Revised Date: 26 March 2018
Accepted Date: 26 March 2018

Please cite this article as: D. Wu, A. Liu, D. Jia, Density functional reactivity theory characterizing the reactivity of frustrated Lewis pairs, *Computational & Theoretical Chemistry* (2018), doi: <https://doi.org/10.1016/j.comptc.2018.03.027>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Density functional reactivity theory characterizing the reactivity of frustrated Lewis pairs

Dongling Wu, Anjie Liu, Dianzeng Jia*

Key Laboratory of Energy Materials Chemistry, Ministry of Education, Key Laboratory of Advanced Functional Materials, Autonomous Region, Institute of Applied Chemistry, Xinjiang University, Urumqi, Xinjiang 830046, China.

*Corresponding author, e-mail: jdz@xju.edu.cn, tel.: +86-991-8581183.

Abstract

Previously, theoretical studies for 2,6-lutidine/ BR_3 ($R = F, Me, C_6F_5$) (J. Phys. Chem. A2010) and $PR_3/B(C_6F_5)_3$ ($R = Me, Ph, tBu, C_6F_5$) (Chem. Phys. Lett.2012) pairs have been carried out by us to fully understand the reactivity differences between classical Lewis adducts and frustrated Lewis pairs (FLPs). Fukui function, as a conceptual DFT-based reactivity descriptor, was calculated and shown to explain well the observed reactivity of these Lewis pairs. The initially promising results encouraged us to focus on the reactivity for a huge number of published FLPs by using other conceptual DFT-based reactivity descriptors. In this work, various descriptors were used to understand the intermolecular and intramolecular Lewis centers' reactivities of FLPs. The calculated results shown that, although there are a few exceptions, the adducts and "frustrated complex" of nonreactive pairs and reactive pairs are characterized by the opposite reactivity descriptors. The reactivity of Lewis pairs evaluated by reactivity descriptors also achieves some thermodynamic support from the Gibbs free energies of H_2 activation.

Keywords: FLPs, chemical reactivity, Conceptual DFT, reactivity descriptors

Download English Version:

<https://daneshyari.com/en/article/7838845>

Download Persian Version:

<https://daneshyari.com/article/7838845>

[Daneshyari.com](https://daneshyari.com)