


Author's Accepted Manuscript

Human serum interactions with phenolic and aroma substances of Kaffir (*Citrus hystrix*) and Key lime (*Citrus aurantifolia*) juices

Rajamohamed Beema Shafreen, Martyna Lubinska, Anna Róžańska, Tomasz Dymerski, Jacek Namieśnik, Elena Katrich, Shela Gorinstein



www.elsevier.com/locate/jlumin

PII: S0022-2313(18)30430-7
DOI: <https://doi.org/10.1016/j.jlumin.2018.04.010>
Reference: LUMIN15525

To appear in: *Journal of Luminescence*

Received date: 8 March 2018
Revised date: 2 April 2018
Accepted date: 5 April 2018

Cite this article as: Rajamohamed Beema Shafreen, Martyna Lubinska, Anna Róžańska, Tomasz Dymerski, Jacek Namieśnik, Elena Katrich and Shela Gorinstein, Human serum interactions with phenolic and aroma substances of Kaffir (*Citrus hystrix*) and Key lime (*Citrus aurantifolia*) juices, *Journal of Luminescence*, <https://doi.org/10.1016/j.jlumin.2018.04.010>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Human serum interactions with phenolic and aroma substances of Kaffir (*Citrus hystrix*) and Key lime (*Citrus aurantifolia*) juices

Rajamohamed Beema Shafreen^{b,*}, Martyna Lubinska^a, Anna Róžańska^a, Tomasz Dymerski^a, Jacek Namieśnik^a, Elena Katrich^c, Shela Gorinstein^{c,*}

^a*Gdańsk University of Technology, Faculty of Chemistry, Department of Analytical Chemistry, Gdańsk 80-233, Poland*

^b*Molecular Nanomedicine Research Unit, Centre for Nanoscience and Nanotechnology, Sathyabama University, Chennai -600119, TN, India*

^c*Institute for Drug Research, School of Pharmacy, Hadassah Medical School, The Hebrew University, Jerusalem 9112001, Israel*

*Corresponding authors:

beema.shafreen@gmail.com

shela.gorin@mail.huji.ac.il

gorin@cc.huji.ac.il

Abstract

To understand the therapeutic application of polyphenols extracted from Kaffir (PolKaf) and Key (PolKey) limes different analytical methods were applied. Based on quantitative analysis by two dimensional gas chromatography (GCxGC) and time of flight mass spectroscopy (TOFMS) it can be observed that the biggest differences in the contents of

Download English Version:

<https://daneshyari.com/en/article/7839906>

Download Persian Version:

<https://daneshyari.com/article/7839906>

[Daneshyari.com](https://daneshyari.com)