


Accepted Manuscript

Two-dimensional ^1H and ^1H -detected NMR study of a heterogeneous biocatalyst using fast MAS at high magnetic fields

Sabu Varghese, Peter J. Halling, Daniel Häussinger, Stephen Wimperis



PII: S0926-2040(18)30013-4

DOI: [10.1016/j.ssnmr.2018.03.003](https://doi.org/10.1016/j.ssnmr.2018.03.003)

Reference: YSNMR 832

To appear in: *Solid State Nuclear Magnetic Resonance*


Received Date: 5 February 2018

Revised Date: 13 March 2018

Accepted Date: 13 March 2018

Please cite this article as: S. Varghese, P.J. Halling, D. Häussinger, S. Wimperis, Two-dimensional ^1H and ^1H -detected NMR study of a heterogeneous biocatalyst using fast MAS at high magnetic fields, *Solid State Nuclear Magnetic Resonance* (2018), doi: 10.1016/j.ssnmr.2018.03.003.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



^1H MAS NMR at high spinning rates and high magnetic fields is used to investigate the structural integrity of an enzyme covalently immobilized on porous silica

Download English Version:

<https://daneshyari.com/en/article/7844509>

Download Persian Version:

<https://daneshyari.com/article/7844509>

[Daneshyari.com](https://daneshyari.com)