


Accepted Manuscript

Graphene-based ordered framework with a diverse range of carbon polygons formed in zeolite nanochannels

Hiroto Nishihara, Hiroyuki Fujimoto, Hiroyuki Itoi, Keita Nomura, Hideki Tanaka, Minoru T. Miyahara, Patrick A. Bonnaud, Ryuji Miura, Ai Suzuki, Naoto Miyamoto, Nozomu Hatakeyama, Akira Miyamoto, Kazutaka Ikeda, Toshiya Otomo, Takashi Kyotani



PII: S0008-6223(17)31288-5

DOI: [10.1016/j.carbon.2017.12.055](https://doi.org/10.1016/j.carbon.2017.12.055)

Reference: CARBON 12685

To appear in: *Carbon*


Received Date: 3 November 2017

Revised Date: 14 December 2017

Accepted Date: 15 December 2017

Please cite this article as: H. Nishihara, H. Fujimoto, H. Itoi, K. Nomura, H. Tanaka, M.T. Miyahara, P.A. Bonnaud, R. Miura, A. Suzuki, N. Miyamoto, N. Hatakeyama, A. Miyamoto, K. Ikeda, T. Otomo, T. Kyotani, Graphene-based ordered framework with a diverse range of carbon polygons formed in zeolite nanochannels, *Carbon* (2018), doi: 10.1016/j.carbon.2017.12.055.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



ACCEPTED MANUSCRIPT

Download English Version:

<https://daneshyari.com/en/article/7849153>

Download Persian Version:

<https://daneshyari.com/article/7849153>

[Daneshyari.com](https://daneshyari.com)