



Original article

Multidetector computed tomography for preoperative identification of left gastric vein location in patients with gastric cancer

KENTARO KAWASAKI, SHINGO KANAJI, IWAO KOBAYASHI, TADATOSHI FUJITA, HIROAKI KOMINAMI, KIMIHIKO UENO, SHINOBU TSUTIDA, MASAKAZU OHNO, MASATO OHSAWA, YASUHIRO FUJINO, MASAHIRO TOMINAGA, and TAKESHI NAKAMURA

Division of Gastroenterological Surgery, Hyogo Cancer Center, 13-70 Kitaouji-cho, Akashi, Hyogo 673-8558, Japan

Abstract

Background. The area near the left gastric vein (LGV) is a challenging site at which to perform dissection of the lymph nodes during gastrectomy. Therefore, knowledge of the precise location of the LGV is important. The objective of this study was to examine the usefulness of multidetector computed tomography (MDCT) for the identification of the LGV.

Methods. Eighty-one patients with gastric cancer underwent MDCT, which was performed with contrast media in 76 patients and without contrast media in 5 patients. A 5-mm thin slice of the frontal image was reconstructed. These images were examined preoperatively to detect the location of the LGV. Upon gastrectomy, the LGV was identified and its location compared to that determined by MDCT.

Results. The LGV was identified by MDCT in 76 of the 81 patients (93.8%). The LGV was subsequently located during the operation in all 81 patients. The LGV was located dorsal to the common hepatic artery in 40 patients (49.4%), ventral to the common hepatic artery in 18 patients (22.2%), ventral to the splenic artery in 17 patients (21.0%), dorsal to the splenic artery in 2 patients (2.5%) and in other positions in 4 patients (4.9%). In all patients, the location of the LGV detected using MDCT was consistent with that identified during gastrectomy. In the 4 patients with relatively unusual locations of the LGV, these 4 LGV variants were identified preoperatively by MDCT.

Conclusion. MDCT was useful for identifying the location of the LGV prior to gastrectomy.

Key words Left gastric vein · Multidetector computed tomography · MDCT · Gastric cancer

Introduction

In Japan, the perigastric lymph nodes are usually dissected in patients with gastric cancer [1, 2]. In our experi-

ence, the area near the left gastric vein (LGV) is one of the most challenging sites at which to perform dissection, because of its various anatomical variants. Therefore, knowledge of the precise location of the LGV preoperatively may make it possible to avoid unnecessary bleeding.

In Japan, computed tomography (CT) scanning from the chest to the abdomen is routinely performed to check for the presence of metastases in patients with gastric cancer. Three-dimensional computed tomographic angiography (3DCTA) is a reliable method for determining the location and anatomy of the vessels around the stomach [3–8]. However, this method requires an additional CT scan over and above the routine CT, so extra cost is incurred. On the other hand, the recent development of multidetector computed tomography (MDCT) now makes it possible to obtain highly detailed information, including high-resolution data on perigastric vascular anatomy.

In the present study, the location of the LGV was assessed during preoperative MDCT used to perform the routine check for metastasis, and these preoperative results were compared to the location of the LGV visualized during the actual operation. The aim of this study was to examine the usefulness of MDCT for identification of the location of the LGV.

Patients and methods

Patients

Eighty-one patients with primary gastric cancer were enrolled in this study. All patients underwent MDCT preoperatively to detect metastasis from the chest to the abdomen. MDCT was performed with contrast media in 76 patients and without contrast media in 5 patients who were allergic to contrast agents. In one patient with arterial and venous variants detected by MDCT,




Fig. 1A–D. Multidetector computed tomography (MDCT) and three-dimensional computed tomographic angiography (3DCTA) imaging in case 31; arterial and venous variants were detected by MDCT (**A** and **B**), so 3DCTA was performed additionally (**C** and **D**). The left gastric vein (LGV; *arrow*) starts from the stomach (**B**) and ends in the hepatic vein (HV; **A**). The venous phase of the 3DCTA imaging showed that the LGV flowed into the HV (**C**). The arterial

phase of the 3DCTA imaging showed that the common hepatic artery (*CHA*) started from the superior mesenteric artery (*SMA*), and the left hepatic artery (*LHA*) started from the left gastric artery (*LGA*; **D**). *SA*, Splenic artery; *CA*, celiac artery; *PV*, portal vein; *SV*, splenic vein; *SMV*, superior mesenteric vein; *RHA*, right hepatic artery; *GDA*, gastroduodenal artery; *HV*, hepatic vein

3DCTA was performed additionally (Fig. 1). All patients underwent gastrectomy, performed at the Gastroenterological Department of Surgery, Hyogo Cancer Center, Hyogo, Japan, between September 2007 and May 2009. Open gastrectomy was performed in 34 patients and laparoscopic gastrectomy in 47 patients.

MDCT and location of LGV

MDCT was performed using an Aquilion 16-row multidetector CT scanner (Toshiba Medical Systems, Tokyo, Japan). Contrast medium (100 ml at a rate of 2 ml/s) was intravenously injected. After 70 s, the patients were scanned from the chest to abdomen. A

5-mm thin slice of the standard frontal image was reconstructed, and these images were examined preoperatively by surgeons (K. K. and S. K.) to detect the location of the LGV. For the purposes of this study, the vessel connecting the portal or splenic vein to the stomach was defined as the LGV (Figs. 2 and 3). The LGV was described as one of the following five types based on its location: (A) dorsal to the common hepatic artery, (B) ventral to the common hepatic artery, (C) ventral to the splenic artery, (D) dorsal to the splenic artery, or (E) other (Fig. 4). The location of the LGV was confirmed during the gastrectomy, and this result was compared to the location determined preoperatively by MDCT imaging.

Download English Version:

<https://daneshyari.com/en/article/7852053>

Download Persian Version:

<https://daneshyari.com/article/7852053>

[Daneshyari.com](https://daneshyari.com)