

Accepted Manuscript

Histidine-rich glycoprotein-induced vascular normalization improves EPR-mediated drug targeting to and into tumors

Benjamin Theek, Maike Baues, Felix Gremse, Robert Pola, Michal Pechar, Inka Negwer, Kaloian Koynov, Benjamin Weber, Matthias Barz, Willi Jahnen-Dechent, Gert Storm, Fabian Kiessling, Twan Lammers



PII: S0168-3659(18)30257-8
DOI: doi:[10.1016/j.jconrel.2018.05.002](https://doi.org/10.1016/j.jconrel.2018.05.002)
Reference: COREL 9284
To appear in: *Journal of Controlled Release*
Received date: 29 January 2018
Revised date: 26 April 2018
Accepted date: 2 May 2018

Please cite this article as: Benjamin Theek, Maike Baues, Felix Gremse, Robert Pola, Michal Pechar, Inka Negwer, Kaloian Koynov, Benjamin Weber, Matthias Barz, Willi Jahnen-Dechent, Gert Storm, Fabian Kiessling, Twan Lammers, Histidine-rich glycoprotein-induced vascular normalization improves EPR-mediated drug targeting to and into tumors. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Corel(2017), doi:[10.1016/j.jconrel.2018.05.002](https://doi.org/10.1016/j.jconrel.2018.05.002)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Histidine-rich glycoprotein-induced vascular normalization improves EPR-mediated drug targeting to and into tumors

Benjamin Theek^{1,2,1}, Maike Baues^{1,1}, Felix Gremse¹, Robert Pola³, Michal Pechar³,
Inka Negwer⁴, Kaloian Koynov⁴, Benjamin Weber⁵, Matthias Barz⁵,
Willi Jahnen-Dechent⁶, Gert Storm^{2,7}, Fabian Kiessling¹, Twan Lammers^{1,2,7,*}
tlammers@ukaachen.de

¹Department of Nanomedicine and Theranostics, Institute for Experimental Molecular Imaging,
RWTH Aachen University Clinic and Helmholtz Institute for Biomedical Engineering, Aachen, Germany

²Department of Targeted Therapeutics, Biomaterial Science and Technology, University of Twente,
Enschede, The Netherlands

³Institute of Macromolecular Chemistry, Academy of Sciences of the Czech Republic, Prague, Czech
Republic

⁴Max Planck Institute for Polymer Research, Mainz, Germany

⁵Institute of Organic Chemistry, Johannes Gutenberg University, Mainz, Germany

⁶Biointerface Laboratory, RWTH Aachen University Clinic and Helmholtz Institute for Biomedical
Engineering, Aachen, Germany

⁷Department of Pharmaceutics, Utrecht Institute for Pharmaceutical Sciences, Utrecht University, Utrecht,
The Netherlands

*Corresponding author.

ABSTRACT: Tumors are characterized by leaky blood vessels, and by an abnormal and heterogeneous vascular network. These pathophysiological characteristics contribute to the enhanced permeability and retention (EPR) effect, which is one of the key rationales for

¹ These authors contributed equally to this study

Download English Version:

<https://daneshyari.com/en/article/7859485>

Download Persian Version:

<https://daneshyari.com/article/7859485>

[Daneshyari.com](https://daneshyari.com)