High-frequency, low-intensity ultrasound and microbubbles enhance nerve blockade

Kathleen Cullion, Claudia M. Santamaria, Changyou Zhan, David Zurakowski, Tao Sun, Nathan J. McDannold, Daniel S. Kohane

PII: S0168-3659(18)30095-6
DOI: doi:10.1016/j.jconrel.2018.02.027
Reference: COREL 9175

To appear in: *Journal of Controlled Release*

Received date: 5 July 2017
Revised date: 5 February 2018
Accepted date: 16 February 2018

Please cite this article as: Kathleen Cullion, Claudia M. Santamaria, Changyou Zhan, David Zurakowski, Tao Sun, Nathan J. McDannold, Daniel S. Kohane , High-frequency, low-intensity ultrasound and microbubbles enhance nerve blockade. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Corel(2018), doi:10.1016/j.jconrel.2018.02.027

High-frequency, low-intensity ultrasound and microbubbles enhance nerve blockade

Kathleen Cullion[1], Claudia M. Santamaria[1], Changyou Zhan[1,2], David Zurakowski[3], Tao Sun[4], Nathan J. McDannold[4], and Daniel S. Kohane[1,3]

1. Laboratory for Biomaterials and Drug Delivery, Division of Critical Care Medicine, Department of Anesthesiology, Boston Children's Hospital, Harvard Medical School, Boston, MA, United States
2. Department of Pharmacology, School of Basic Medical Sciences, Fudan University & Key Laboratory of Smart Drug Delivery (Fudan University), Ministry of Education, Shanghai, China
3. Department of Anesthesiology, Boston Children's Hospital, Harvard Medical School, Boston, MA, United States
4. Focused Ultrasound Laboratory, Department of Radiology, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, United States

Correspondence:

Daniel S. Kohane, M.D., Ph.D.

300 Longwood Ave

Enders Building, Room 361

Boston Children's Hospital

Boston, MA 02115

Email: Daniel.Kohane@childrens.harvard.edu

Introduction