

Accepted Manuscript

Cardiac progenitor cells activated by mitochondrial delivery of resveratrol enhance the survival of a doxorubicin-induced cardiomyopathy mouse model via the mitochondrial activation of a damaged myocardium

Jiro Abe, Yuma Yamada, Takeda Atsuhito, Hideyoshi Harashima

PII: S0168-3659(17)31020-9
DOI: doi:[10.1016/j.jconrel.2017.11.024](https://doi.org/10.1016/j.jconrel.2017.11.024)
Reference: COREL 9054

To appear in: *Journal of Controlled Release*

Received date: 25 July 2017
Revised date: 24 October 2017
Accepted date: 13 November 2017

Please cite this article as: Jiro Abe, Yuma Yamada, Takeda Atsuhito, Hideyoshi Harashima, Cardiac progenitor cells activated by mitochondrial delivery of resveratrol enhance the survival of a doxorubicin-induced cardiomyopathy mouse model via the mitochondrial activation of a damaged myocardium. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. *Corel*(2017), doi:[10.1016/j.jconrel.2017.11.024](https://doi.org/10.1016/j.jconrel.2017.11.024)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Cardiac progenitor cells activated by mitochondrial delivery of resveratrol enhance the survival of a doxorubicin-induced cardiomyopathy mouse model via the mitochondrial activation of a damaged myocardium

Jiro Abe^{1,3}, Yuma Yamada^{2,3}, Takeda Atsuhito¹, Hideyoshi Harashima^{2,*}

¹Department of Pediatrics, Graduate School of Medicine, Hokkaido University, Kita-15, Nishi 7, Kita-ku, Sapporo 060-8638, Japan

²Laboratory for molecular design of pharmaceuticals, Faculty of Pharmaceutical Sciences, Hokkaido University, Kita-12, Nishi-6, Kita-ku, Sapporo 060-0812, Japan

³These authors equally contributed to this study.

*Corresponding author: Laboratory for molecular design of pharmaceuticals, Faculty of Pharmaceutical Sciences, Hokkaido University, Kita-12, Nishi-6, Kita-ku, Sapporo 060-0812, Japan

Tel: +81-11-706-3919 Fax: +81-11-706-4879

E-mail: harasima@pharm.hokudai.ac.jp

Download English Version:

<https://daneshyari.com/en/article/7860662>

Download Persian Version:

<https://daneshyari.com/article/7860662>

[Daneshyari.com](https://daneshyari.com)