Enhanced FcRn-dependent transepithelial delivery of IgG by Fc-engineering and polymerization

Stian Foss, Algirdas Grevys, Kine Marita Knudsen Sand, Malin Bern, Pat Blundell, Terje E. Michaelsen, Richard J. Pleass, Inger Sandlie, Jan Terje Andersen

PII: S0168-3659(15)30284-4
DOI: doi: 10.1016/j.jconrel.2015.12.033
Reference: COREL 8032

To appear in: *Journal of Controlled Release*

Received date: 24 September 2015
Revised date: 14 December 2015
Accepted date: 19 December 2015

Please cite this article as: Stian Foss, Algirdas Grevys, Kine Marita Knudsen Sand, Malin Bern, Pat Blundell, Terje E. Michaelsen, Richard J. Pleass, Inger Sandlie, Jan Terje Andersen, Enhanced FcRn-dependent transepithelial delivery of IgG by Fc-engineering and polymerization, *Journal of Controlled Release* (2015), doi: 10.1016/j.jconrel.2015.12.033

Enhanced FcRn-dependent transepithelial delivery of IgG by Fc-engineering and polymerization

Stian Foss[1,2], Algirdas Grevys[1,2], Kine Marita Knudsen Sand[1,2], Malin Bern[1,2], Pat Blundell[5], Terje E. Michaelsen[3,4], Richard J. Pleass[5], Inger Sandlie[1,2] and Jan Terje Andersen[1,2]

[1]Centre for Immune Regulation (CIR) and Department of Biosciences, University of Oslo, N-0316 Oslo, Norway. [2]Department of Immunology and CIR, Oslo University Hospital, Rikshospitalet and University of Oslo, N-0372 Oslo, Norway. [3]Department of Bacteriology and Immunology, Norwegian Institute of Public Health, Oslo, Norway. [4]Department of Chemical Pharmacy, School of Pharmacy, University of Oslo, Oslo, Norway. [5]Liverpool School of Tropical Medicine, Pembroke Place, Liverpool, L3 5QA, UK.

To whom correspondence should be addressed: Jan Terje Andersen, Centre for Immune Regulation (CIR) and Department of Immunology, Oslo University Hospital, Rikshospitalet and University of Oslo, PO Box 4956, Oslo N-0424, Norway. E-mail: j.t.andersen@medisin.uio.no.

Keywords: FcRn, pH-dependent binding, IgG, Fc-fusion, transcytosis, epithelial, mucosal barrier.