Layered double hydroxides as effective carrier for anticancer drugs and tailoring of release rate through interlayer anions

Sudipta Senapati, Ravi Thakur, Shiv Prakash Verma, Shivali Duggal, Durga Prasad Mishra, Parimal Das, T. Shripathi, Mohan Kumar, Dipak Rana, Pralay Maiti

PII: S0168-3659(16)30013-X
DOI: doi: 10.1016/j.jconrel.2016.01.016
Reference: COREL 8072

To appear in: *Journal of Controlled Release*

Received date: 16 September 2015
Revised date: 28 December 2015
Accepted date: 11 January 2016

Please cite this article as: Sudipta Senapati, Ravi Thakur, Shiv Prakash Verma, Shivali Duggal, Durga Prasad Mishra, Parimal Das, T. Shripathi, Mohan Kumar, Dipak Rana, Pralay Maiti, Layered double hydroxides as effective carrier for anticancer drugs and tailoring of release rate through interlayer anions, *Journal of Controlled Release* (2016), doi: 10.1016/j.jconrel.2016.01.016

Layered Double Hydroxides as Effective Carrier for Anticancer Drugs and Tailoring of Release Rate through Interlayer Anions

Sudipta Senapati,[1] Ravi Thakur,[2] Shiv Prakash Verma,[3] Shivali Duggal,[2] Durga Prasad Mishra,[2] Parimal Das,[3] T. Shripathi,[4] Mohan Kumar,[5] Dipak Rana,[6] and Pralay Maiti[*1]

[1]School of Materials Science and Technology, Indian Institute of Technology (Banaras Hindu University), Varanasi 221 005, India

[2]Cell Death Research Laboratory, Division of Endocrinology, CSIR-Central Drug Research Institute, Lucknow 226031, India

[3]Centre for Genetic Disorders, Faculty of Science, Banaras Hindu University, Varanasi 221005, India

[4]UGC-DAE CSR, University Campus, Khandwa Road, Indore, 452 001, India

[5]Department of Pathology, Institute of Medical Science, Banaras Hindu University, Varanasi 221005, India

[6]Industrial Membrane Research Institute, Department of Chemical and Biological Engineering, University of Ottawa, 161 Louis Pasteur St., Ottawa, ON, Canada KIN 6N5

[*]Correspondence should be made to Pralay Maiti (pmaiti.mst@itbhu.ac.in)