## **Accepted Manuscript**

Clifford algebra valued boundary integral equations for three-dimensional elasticity

Li-Wei Liu, Hong-Ki Hong

PII: \$0307-904X(17)30583-8 DOI: 10.1016/j.apm.2017.09.031

Reference: APM 11976

To appear in: Applied Mathematical Modelling

Received date: 2 January 2016 Revised date: 30 June 2017

Accepted date: 18 September 2017



Please cite this article as: Li-Wei Liu, Hong-Ki Hong, Clifford algebra valued boundary integral equations for three-dimensional elasticity, *Applied Mathematical Modelling* (2017), doi: 10.1016/j.apm.2017.09.031

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

#### ACCEPTED MANUSCRIPT

#### Highlights

- Clifford analysis is applied to solve 3D problems of elasticity.
- $\bullet$  BIEs for Clifford algebra valued harmonic functions are derived.
- The singularity-free BIEs are constructed without extra equations/variables/information.
- The Clifford algebra valued singularity-free BEM is developed by a regular quadrature.
- 3D problems of elasticity can be solved using the Clifford algebra valued RFM

### Download English Version:

# https://daneshyari.com/en/article/8052093

Download Persian Version:

https://daneshyari.com/article/8052093

<u>Daneshyari.com</u>