
## Accepted Manuscript

Effects of different covering systems and carbon nitrogen ratios on nitrogen removal in surface flow constructed wetlands



PII:	S0959-6526(17)32477-0
DOI:	10.1016/j.jclepro.2017.10.170
Reference:	JCLP 10962
To appear in:	Journal of Cleaner Production
Received Date:	21 June 2017
Revised Date:	30 September 2017
Accepted Date:	15 October 2017

Please cite this article as: Xiaowen Ding, Ying Xue, Yong Zhao, Weihua Xiao, Yuan Liu, Jiagang Liu, Effects of different covering systems and carbon nitrogen ratios on nitrogen removal in surface flow constructed wetlands, *Journal of Cleaner Production* (2017), doi: 10.1016/j.jclepro. 2017.10.170

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



- 1 Amount of words: 8398
- 2 Effects of different covering systems and carbon nitrogen ratios on nitrogen
  3 removal in surface flow constructed wetlands
- 4 Xiaowen Ding<sup>a,\*</sup>, Ying Xue<sup>a</sup>, Yong Zhao<sup>b</sup>, Weihua Xiao<sup>b</sup>, Yuan Liu<sup>a</sup>, Jiagang Liu<sup>c</sup>
- <sup>5</sup> <sup>a</sup>Key Laboratory of Regional Energy and Environmental Systems Optimization, Ministry
- 6 of Education, North China Electric Power University, Beijing, 102206, China
- <sup>7</sup> <sup>b</sup>State Key Laboratory of Simulation and Regulation of Water Cycle in River Basin,
- 8 China Institute of Water Resources and Hydropower Research, Beijing 100038, China
- 9 °Tsinghua Holdings Human Settlements Environment Institute, Beijing, 100083, China
- 10

## 11 Highlights:

- 12 Effects of different covering systems and COD/N on nitrogen removal were studied.
- 13 Nitrogen was divided into ORN,  $NH_4^+$ -N and  $NO_3^-$ -N.
- Plants favored ORN and  $NH_4^+$ -N removal, and large coverage favored  $NO_3^-$ -N removal.
- 15 Nitrification was the key step in nitrogen removal.
- 16 Results provide a theoretical basis for non-point source pollution control using SFW.

\* Corresponding author. North China Electric Power University, 2 Beinong Road, Changping District, Beijing, 102206, China

*E-mail address:* <u>binger2000dxw@hotmail.com</u> (X. Ding), <u>astridxuey@163.com</u> (Y. Xue), <u>zhaoyong@iwhr.com</u> (Y. Zhao), <u>xwsen998@163.com</u> (W. Xiao), <u>ly921105@gmail.com</u> (Y. Liu), <u>liujiagang@139.com</u> (J. Liu).

Download English Version:

## https://daneshyari.com/en/article/8099937

Download Persian Version:

https://daneshyari.com/article/8099937

Daneshyari.com