

Accepted Manuscript

Efficiency of BRDF sampling and bias on the average photometric behavior

Frédéric Schmidt, Sébastien Bourguignon


PII: S0019-1035(17)30110-0
DOI: [10.1016/j.icarus.2018.06.025](https://doi.org/10.1016/j.icarus.2018.06.025)
Reference: YICAR 12946

To appear in: *Icarus*

Received date: 6 February 2017
Revised date: 11 June 2018
Accepted date: 25 June 2018

Please cite this article as: Frédéric Schmidt, Sébastien Bourguignon, Efficiency of BRDF sampling and bias on the average photometric behavior, *Icarus* (2018), doi: [10.1016/j.icarus.2018.06.025](https://doi.org/10.1016/j.icarus.2018.06.025)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Highlights

- Definition of the efficiency of a set of geometries to constrain Hapke photometric parameters
- Best efficiency is principal plane with high incidence
- Study of the heterogeneity in dataset
- Separability test of chi-square is able to detect separability for some parameters but not for all.

ACCEPTED MANUSCRIPT

Download English Version:

<https://daneshyari.com/en/article/8133623>

Download Persian Version:

<https://daneshyari.com/article/8133623>

[Daneshyari.com](https://daneshyari.com)