

New polarimetric and spectroscopic evidence of anomalous enrichment in spinel-bearing Calcium-Aluminium-rich Inclusions among L-type asteroids

M. Devogèle, P. Tanga, A. Cellino, Ph. Bendjoya, J.-P. Rivet, J. Surdej, D. Vernet, J.M. Sunshine, S.J. Bus, L. Abe, S. Bagnulo, G. Borisov, H. Campins, B. Carry, J. Licandro, W. McLean, N. Pinilla-Alonso

PII: S0019-1035(17)30027-1
DOI: 10.1016/j.icarus.2017.12.026
Reference: YICAR 12746

To appear in: *Icarus*

Received date: 15 January 2017
Revised date: 25 November 2017
Accepted date: 18 December 2017

Please cite this article as: M. Devogèle, P. Tanga, A. Cellino, Ph. Bendjoya, J.-P. Rivet, J. Surdej, D. Vernet, J.M. Sunshine, S.J. Bus, L. Abe, S. Bagnulo, G. Borisov, H. Campins, B. Carry, J. Licandro, W. McLean, N. Pinilla-Alonso, New polarimetric and spectroscopic evidence of anomalous enrichment in spinel-bearing Calcium-Aluminium-rich Inclusions among L-type asteroids, *Icarus* (2017), doi: 10.1016/j.icarus.2017.12.026

Highlights

- Spectroscopic and polarimetric observation of Barbarian/L-type asteroids
- Hapke modeling of asteroid spectra including model of space-weathering
- CAIs were found to be the most probable of the large inversion angle of Barbarians