

Heavy flavor and vector boson associated production

Carlotta Favaro *on behalf of the CMS Collaboration*

Universität Zürich, Physik-Institut, Winterthurerstr. 190, 8057 Zürich, Switzerland

Abstract

The mechanism of production of heavy-flavoured mesons, containing b or c quarks, in association with vector bosons, W or Z/γ^* , in the Standard Model is only partially understood. The study of events with one or two well-identified and isolated leptons accompanied by b-jets or secondary vertices is therefore crucial to refine the theoretical calculations in perturbative QCD, as well as validate associated Monte Carlo techniques. The deep understanding of these processes is furthermore required by Higgs and BSM analyses with similar final states. Using the LHC proton-proton collision data collected in 2010 and 2011 at a centre of mass energy of 7 TeV by the CMS detector, preliminary measurements of the $Z/\gamma^* + b(b)$ cross sections and angular correlations are presented. Finally, the study of the $W + c$ production rate with respect to the W charge and $W + \text{light jets}$ rates allows to probe the strange quark content of the proton. These results are also presented.

Keywords: LHC, CMS, Z, W, bottom, charm

1. Measurement of Z and heavy flavor associated production

1.1. Motivations

The associated production of a Z/γ^* boson and b-quarks is of great relevance at the Large Hadron Collider. It represents a significant background to the search of the Standard Model Higgs boson produced in association with a vector boson and decaying into a $b\bar{b}$ pair, and it's a possible discovery channel for models with an extended Higgs sector, such as the two Higgs-doublet model.

Besides, the $Z/\gamma^* + \text{heavy flavor}$ production is a good test of perturbative QCD and of Monte Carlo simulation. The description of the dynamics of b-quarks remains problematic, especially in the soft/collinear region, dominated by the gluon-splitting contribution. Two different approaches are implemented in the simulation of the $Z/\gamma^* + b$ production: the four-flavor, which excludes the b-quarks from the proton parton density functions, and the five-flavor, which instead integrates the b-quarks in the PDFs. The measurement of the inclusive $Z/\gamma^* + b$ -jet cross-section, and of the angular correlations in $Z/\gamma^* + b\bar{b}$ events can help identifying the modeling that provides the best description of data.

1.2. $Z/\gamma^* + b$ and $Z/\gamma^* + b\bar{b}$ cross-section measurements

The inclusive cross-section of the Z/γ^* produced in association with at least one b-jet [1] as well as the inclusive $Z/\gamma^* + b\bar{b}$ production cross-section [2] are measured in a data sample corresponding to an integrated luminosity of 2.2 fb^{-1} , collected in 2011 with the CMS detector [3].

The Z/γ^* boson is reconstructed in its leptonic decays $Z/\gamma^* \rightarrow \mu\mu$ and $Z/\gamma^* \rightarrow ee$. The event sample is thus selected by requiring the presence of two isolated muons (electrons) with $p_T > 20$

GeV (25 GeV) and $|\eta| < 2.1$ (2.5). Only dilepton pairs fulfilling the condition $60 < M(l\bar{l}) < 120 \text{ GeV}$ for the $Z/\gamma^* + b$ -jet cross-section measurement, and by $76 < M(l\bar{l}) < 106 \text{ GeV}$ for the $Z/\gamma^* + b\bar{b}$ analysis, are retained. Jets are reconstructed from particle-flow objects [8] clustered with the anti- k_T algorithm [9], and are required to have $p_T > 25 \text{ GeV}$, $|\eta| < 2.1$, and to be separated from each of the leptons by $\Delta R > 0.5$. The b fraction is enhanced by applying the Simple Secondary Vertex (SSV) b-tagging algorithm [10], whose discriminator is a function of the distance between the primary interaction and the secondary vertex from the heavy flavor decay. The algorithm is applied in its High Purity working point, that requires at least three tracks associated to the vertex, in the $Z/\gamma^* + b$ cross-section estimation, and in its High Efficiency working point, requiring at least two tracks, in the $Z/\gamma^* + b\bar{b}$ measurement.

The main backgrounds to the $Z/\gamma^* + b$ -jet signal are $Z/\gamma^* + \text{light quark/gluon}$ and charm jets and top pair production. A maximum-likelihood fit of the distribution of the secondary vertex mass for the leading b-jet is used to estimate the b-jet purity \mathcal{P} , while the fraction of top pair events $f_{t\bar{t}}$ is extracted by extrapolating into the signal region the number of observed events in the Z upper sideband. The $Z/\gamma^* + b$ -jet cross-section is computed from the number of selected events $N(l\bar{l} + b)$, taking into account the b-tagging efficiency ϵ_b , the lepton acceptance \mathcal{A}_l and selection efficiency ϵ_l , the correction factor C_{hadron} for detector and reconstruction effects, and the integrated luminosity \mathcal{L} , as in the formula

$$\sigma(Z/\gamma^* + b) = \frac{N(l\bar{l} + b) \cdot (\mathcal{P} - f_{t\bar{t}})}{\mathcal{A}_l \cdot C_{hadron} \cdot \epsilon_l \cdot \epsilon_b \cdot \mathcal{L}} \quad (1)$$

The b-tagging and lepton selection efficiency are estimated from Monte Carlo simulation, and rescaled to reproduce the efficiencies measured on data with the tag-and-probe technique. The




Figure 1: Secondary vertex invariant mass distribution for the leading b-jet in events with two leptons in the invariant mass range $76 < M(ll) < 106$ GeV and two b-jets. A maximum-likelihood fit of the vertex mass distribution is used to extract the b-jet purity \mathcal{P} .

correction for detector effects is instead evaluated on the simulation only.

The main systematic uncertainties are related to the b-tagging efficiency (10%) and b-jet purity (5%). The result is $\sigma(Z/\gamma^* + b) = 5.84 \pm 0.08(stat.) \pm 0.72(syst.)^{+0.25}_{-0.55}(theory)$ pb, referred to a phase space defined for the leptons by $60 < M(ll) < 120$ GeV, and for the jets by $p_T > 25$ GeV and $|\eta| < 2.1$. The result is 30% higher than the hadron-level-corrected NLO prediction of $\sigma(Z/\gamma^* + b) = 3.97 \pm 0.47$ pb by MCFM.

The procedure to extract the $Z/\gamma^* + bb$ inclusive cross-section is similar to the one described above. The requirement of the presence of two b-jets suppresses the contamination from $Z/\gamma^* +$ light quark/gluon and charm jets and increases the b-jet purity of the sample. The latter is estimated through a simultaneous maximum-likelihood fit of secondary vertex mass distribution the leading and sub-leading b-tagged jets, as shown in Fig.1, and is estimated to be about 84%. In order to reduce the top background, a cut is applied on the missing transverse energy $\cancel{E}_T < 50$ GeV. The remaining top fraction, about 20%, is evaluated through a maximum-likelihood fit of the dilepton invariant mass distribution. A small contamination from diboson production is instead estimated from Monte Carlo simulation. The hadron-level cross-section of $Z/\gamma^* + bb$ is extracted by applying a set of unfolding matrices, accounting for bin-to-bin migrations in the b-jet multiplicity due to dilepton efficiency and acceptance, b-tagging efficiency, and detector and reconstruction effects. The result is $\sigma(Z/\gamma^* + bb) = 0.37 \pm 0.02(stat.) \pm 0.07(syst.) \pm 0.11(theory)$ pb, in good agreement with the tree-level prediction of $\sigma(Z/\gamma^* + bb) = 0.33 \pm 0.01$ pb by MADGRAPH [6].

Some discrepancies are though observed in the event kinematics, as shown in Fig. 2.





Figure 2: Transverse momentum of the b-jet pair (top) and transverse angular separation between the Z/γ^* and the b-jet pair (bottom), in events with two b-jets. The yellow band in the bottom part represents the statistical uncertainty on the Monte Carlo yields.

1.3. Angular correlation between B-hadrons produced in association with a Z boson

The normalized differential cross-section of $Z/\gamma^* + BB$ -hadrons as function of the angular separation between the B-hadrons, defined by the variable $\Delta R = \sqrt{(\Delta\phi)^2 + (\Delta\eta)^2}$, is measured using a data sample corresponding to 4.6 pb^{-1} collected in 2011 [4].

A sample of Z/γ^* events is selected using the same criteria as for the $Z/\gamma^* + b$ -jet cross-section measurement. The identification of the B-hadrons is performed through the reconstruction of the displaced secondary vertex from their decay, with no use of jets, providing unprecedented sensitivity to B pair production at small angular separation. The measurement refers to the kinematic region defined by two B-hadrons with $p_T > 15$ GeV and $|\eta| < 2$ and two oppositely charged leptons with $p_T > 20$ GeV, $|\eta| < 2.4$ and invariant mass $60 < M(ll) < 120$ GeV.

Download English Version:

<https://daneshyari.com/en/article/8185415>

Download Persian Version:

<https://daneshyari.com/article/8185415>

[Daneshyari.com](https://daneshyari.com)