


Accepted Manuscript

A copper-deficient form of mutant cu/Zn-superoxide dismutase as an early pathological species in amyotrophic lateral sclerosis

Eiichi Tokuda, Takao Nomura, Shinji Ohara, Seiji Watanabe, Koji Yamanaka, Yuta Morisaki, Hidemi Misawa, Yoshiaki Furukawa



PII: S0925-4439(18)30095-4
DOI: doi:[10.1016/j.bbadis.2018.03.015](https://doi.org/10.1016/j.bbadis.2018.03.015)
Reference: BBADIS 65086

To appear in:

Received date: 7 January 2018
Revised date: 27 February 2018
Accepted date: 14 March 2018

Please cite this article as: Eiichi Tokuda, Takao Nomura, Shinji Ohara, Seiji Watanabe, Koji Yamanaka, Yuta Morisaki, Hidemi Misawa, Yoshiaki Furukawa , A copper-deficient form of mutant cu/Zn-superoxide dismutase as an early pathological species in amyotrophic lateral sclerosis. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Bbadis(2018), doi:[10.1016/j.bbadis.2018.03.015](https://doi.org/10.1016/j.bbadis.2018.03.015)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Revision: BBADIS-18-20

A copper-deficient form of mutant Cu/Zn-superoxide dismutase as an early pathological species in amyotrophic lateral sclerosis

Eiichi Tokuda¹⁾, Takao Nomura¹⁾, Shinji Ohara²⁾, Seiji Watanabe³⁾, Koji Yamanaka³⁾, Yuta Morisaki⁴⁾, Hidemi Misawa⁴⁾, and *Yoshiaki Furukawa¹⁾

1) Laboratory for Mechanistic Chemistry of Biomolecules, Department of Chemistry, Keio University, Yokohama 223-8522, JAPAN. 2) Department of Neurology, Matsumoto Medical Center, Matsumoto 399-0021, JAPAN. 3) Department of Neuroscience and Pathobiology, Research Institute of Environmental Medicine, Nagoya University, Nagoya 464-8601, JAPAN. 4) Division of Pharmacology, Faculty of Pharmacy, Keio University, Tokyo 105-8512, JAPAN.

Eiichi Tokuda	eiichi.tokuda@gmail.com
Takao Nomura	nomu@pharm.hokudai.ac.jp
Shinji Ohara	oharas@hosp.go.jp
Seiji Watanabe	swtnb@riem.nagoya-u.ac.jp
Koji Yamanaka	kojiyama@riem.nagoya-u.ac.jp
Yuta Morisaki	yutamrsk@gmail.com
Hidemi Misawa	misawa-hd@pha.keio.ac.jp
Yoshiaki Furukawa	furukawa@chem.keio.ac.jp

*Correspondence should be addressed to

Prof. Yoshiaki Furukawa

Department of Chemistry, Keio University, 3-14-1 Hiyoshi, Kohoku, Yokohama, Kanagawa 223-8522, JAPAN. Tel, +81-45-566-1807; Fax, +81-45-566-1697; E-mail, furukawa@chem.keio.ac.jp.

Download English Version:

<https://daneshyari.com/en/article/8258435>

Download Persian Version:

<https://daneshyari.com/article/8258435>

[Daneshyari.com](https://daneshyari.com)