

Induction of cytochrome P450 4A14 contributes to angiotensin II-induced renal fibrosis in mice

Yunfeng Zhou, Jingwei Yu, Jia Liu, Rong Cao, Wen Su, Sha Li, Shiqi Ye, Chenggang Zhu, Xiaolin Zhang, Hu Xu, Hua Chen, Xiaoyan Zhang, Youfei Guan

PII: S0925-4439(17)30482-9
DOI: https://doi.org/10.1016/j.bbadis.2017.12.028
Reference: BBADIS 65004

To appear in:

Received date: 15 August 2017
Revised date: 15 December 2017
Accepted date: 18 December 2017

Please cite this article as: Yunfeng Zhou, Jingwei Yu, Jia Liu, Rong Cao, Wen Su, Sha Li, Shiqi Ye, Chenggang Zhu, Xiaolin Zhang, Hu Xu, Hua Chen, Xiaoyan Zhang, Youfei Guan , Induction of cytochrome P450 4A14 contributes to angiotensin II-induced renal fibrosis in mice. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Bbadis(2017), https://doi.org/10.1016/j.bbadis.2017.12.028

Induction of Cytochrome P450 4A14 Contributes to Angiotensin II-induced Renal Fibrosis in Mice

Yunfeng Zhou[a], Jingwei Yu[a], Jia Liu[a], Rong Cao[a,b], Wen Su[a], Sha Li[c], Shiqi Ye[a], Chenggang Zhu[d], Xiaolin Zhang[d], Hu Xu[e], Hua Chen[e], Xiaoyan Zhang[e,*], and Youfei Guan[a,e,*]

[a]*AstraZeneca-Shenzhen University Joint Institute of Nephrology, Shenzhen University Health Science Center, Shenzhen 518060, China*;

[b]*Department of Nephrology, the First Affiliated Hospital of Shenzhen University, Shenzhen 518039, China;*

[c]*Department of Pathophysiology, Medical College of Hebei University of Engineering, Handan 056002, China*;

[d]*Asia & Emerging Markets Innovative Medicines, AstraZeneca R&D, Shanghai 201203, China*;

[e]*Advanced Institute of Medical Sciences (AIMS), Dalian Medical University, Dalian 116044, China*

[*]Correspondence and offprint requests to:

Youfei Guan, E-mail: guanyf@dmu.edu.cn; or Xiaoyan Zhang, wserien@163.com

Running title: Role of CYP4A14 in AngII-associated renal fibrosis