

N-glycosylation-negative catalase: a useful tool for exploring the role of hydrogen peroxide in the endoplasmic reticulum

S. Lortz, S. Lenzen, I. Mehmeti

www.elsevier.com/locate/freerad-biomed

PII: S0891-5849(14)01402-6
DOI: http://dx.doi.org/10.1016/j.freeradbiomed.2014.11.024
Reference: FRB12246

To appear in: *Free Radical Biology and Medicine*

Received date: 30 July 2014
Revised date: 7 November 2014
Accepted date: 27 November 2014

Cite this article as: S. Lortz, S. Lenzen, I. Mehmeti, N-glycosylation-negative catalase: a useful tool for exploring the role of hydrogen peroxide in the endoplasmic reticulum, *Free Radical Biology and Medicine,* http://dx.doi.org/10.1016/j.freeradbiomed.2014.11.024

N-glycosylation-negative catalase: a useful tool for exploring the role of hydrogen peroxide in the endoplasmic reticulum

S. Lortz*, S. Lenzen, I. Mehmeti

Institute of Clinical Biochemistry, Hannover Medical School, 30623 Hannover, Germany

*****Address correspondence to:

Dr. Stephan Lortz
Institute of Clinical Biochemistry
Hannover Medical School
30623 Hannover
Germany

Telephone: + 49/511/5323765
Fax: + 49/511/5323584
Lortz.Stephan@mh-hannover.de

Keywords: H_2O_2, catalase; endoplasmic reticulum; disulfide bond formation; protein folding.