


## Accepted Manuscript

Hyperglycemia predicts unfavorable outcomes in acute ischemic stroke patients treated with intravenous thrombolysis among a Chinese population: A prospective cohort study

Sheng-Feng Lin, A-Ching Chao, Han-Hwa Hu, Ruey-Tay Lin, Chih-Hung Chen, Lung Chan, Huey-Juan Lin, Yu Sun, Yung-Yang Lin, Po-Lin Chen, Shinn-Kuang Lin, Ming-Hui Sun, Cheng-Yu Wei, Yu-Te Lin, Jiunn-Tay Lee, Chi-Huey Bai, Taiwan Thrombolytic Therapy for Acute Ischemic Stroke (TTT-AIS) Study Group



PII: S0022-510X(18)30147-3  
DOI: doi:[10.1016/j.jns.2018.03.022](https://doi.org/10.1016/j.jns.2018.03.022)  
Reference: JNS 15837

To appear in: *Journal of the Neurological Sciences*

Received date: 16 January 2018  
Revised date: 22 February 2018  
Accepted date: 13 March 2018

Please cite this article as: Sheng-Feng Lin, A-Ching Chao, Han-Hwa Hu, Ruey-Tay Lin, Chih-Hung Chen, Lung Chan, Huey-Juan Lin, Yu Sun, Yung-Yang Lin, Po-Lin Chen, Shinn-Kuang Lin, Ming-Hui Sun, Cheng-Yu Wei, Yu-Te Lin, Jiunn-Tay Lee, Chi-Huey Bai, Taiwan Thrombolytic Therapy for Acute Ischemic Stroke (TTT-AIS) Study Group, Hyperglycemia predicts unfavorable outcomes in acute ischemic stroke patients treated with intravenous thrombolysis among a Chinese population: A prospective cohort study. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. *Jns*(2017), doi:[10.1016/j.jns.2018.03.022](https://doi.org/10.1016/j.jns.2018.03.022)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Hyperglycemia Predicts Unfavorable Outcomes in Acute Ischemic Stroke Patients Treated with Intravenous Thrombolysis among a Chinese Population: a Prospective Cohort Study

### Authors and Affiliations

Sheng-Feng Lin<sup>a</sup>, MD, A-Ching Chao<sup>b, c</sup>, MD, PhD; Han-Hwa Hu<sup>d, e, f</sup>, MD; Ruey-Tay Lin<sup>b, c</sup>, MD; Chih-Hung Chen<sup>g, h</sup>, MD; Lung Chan<sup>f</sup>, MD, PhD; Huey-Juan Lin<sup>i</sup>, MD, MPH; Yu Sun<sup>j</sup>, MD, PhD; Yung-Yang Lin<sup>k</sup>, MD, PhD; Po-Lin Chen<sup>l</sup>, MD; Shinn-Kuang Lin<sup>m</sup>, MD; Ming-Hui Sun<sup>n</sup>, MD; Cheng-Yu Wei<sup>o</sup>, MD; Yu-Te Lin<sup>p</sup>, MD, PhD; Jiunn-Tay Lee<sup>q</sup>, MD; Chi-Huey Bai<sup>a, r</sup>, PhD (Corresponding Author); on behalf of the Taiwan Thrombolytic Therapy for Acute Ischemic Stroke (TTT-AIS) Study Group

<sup>a</sup>School of Public Health, College of Public Health, Taipei Medical University, Taipei, Taiwan

<sup>b</sup>Graduate Institute of Clinical Medicine and Department of Neurology, College of Medicine, Kaohsiung Medical University, Kaohsiung, Taiwan

<sup>c</sup>Department of Neurology, Kaohsiung Medical University Hospital, Kaohsiung, Taiwan

<sup>d</sup>Graduate Institute of Clinical Medicine, College of Medicine, Taipei Medical University, Taipei, Taiwan

<sup>e</sup>Research Center of Cerebrovascular Disease Treatment, College of Medicine, Taipei Medical University, Taipei, Taiwan

<sup>f</sup>Department of Neurology, Taipei Medical University-Shaung Ho Hospital, Taipei, Taiwan

<sup>g</sup>Department of Neurology, National Cheng Kung University Hospital, Tainan, Taiwan

<sup>h</sup>Department of Neurology, National Cheng Kung University, Tainan, Taiwan

<sup>i</sup>Department of Neurology, Chi Mei Medical Center, Tainan, Taiwan

Download English Version:

<https://daneshyari.com/en/article/8272763>

Download Persian Version:

<https://daneshyari.com/article/8272763>

[Daneshyari.com](https://daneshyari.com)