


Accepted Manuscript

Low adherence to antidepressants is associated with increased mortality in Parkinson disease patients

Gal Shoval, Brendon Stubbs, Ran D. Balicer, Becca Feldman, Moshe Hoshen, Gil Zalsman, Roi Sagy, Eldar Hochman, Abraham Weizman, Amir Krivoy



PII: S1353-8020(17)30279-1

DOI: [10.1016/j.parkreldis.2017.07.032](https://doi.org/10.1016/j.parkreldis.2017.07.032)

Reference: PRD 3377

To appear in: *Parkinsonism and Related Disorders*

Received Date: 4 May 2017

Revised Date: 14 June 2017

Accepted Date: 31 July 2017

Please cite this article as: Shoval G, Stubbs B, Balicer RD, Feldman B, Hoshen M, Zalsman G, Sagy R, Hochman E, Weizman A, Krivoy A, Low adherence to antidepressants is associated with increased mortality in Parkinson disease patients, *Parkinsonism and Related Disorders* (2017), doi: 10.1016/j.parkreldis.2017.07.032.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Shoval et al: Adherence to antidepressants in Parkinson patients and mortality

FULL-LENGTH ARTICLE:

Low adherence to antidepressants is associated with increased mortality in Parkinson disease patients

Gal Shoval^{1,2,3}, Brendon Stubbs^{4,5}, Ran D. Balicer^{1,6}, Becca Feldman¹, Moshe Hoshen¹,
Gil Zalsman^{2,3,7}, Roi Sagy^{1,3}, Eldar Hochman^{2,3}, Abraham Weizman^{2,3,8}, Amir Krivoy^{1,2,3}

- 1 Clalit Research Institute, Chief Physician Office, Clalit Health Services, Tel Aviv, Israel
- 2 Geha Mental Health Center, Petah Tiqva, Israel
- 3 Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel
- 4 Physiotherapy Department, South London and Maudsley NHS Foundation Trust, Denmark Hill, London SE5 8AZ, United Kingdom;
- 5 Health Service and Population Research Department, Institute of Psychiatry, Psychology and Neuroscience, King's College London, De Crespigny Park, London Box SE5 8AF, United Kingdom
- 6 Public Health Department, Faculty of Health Sciences, Ben-Gurion University, Beer-Sheva, Israel
- 7 Division of Molecular Imaging and Neuropathology, Department of Psychiatry, Columbia University, New York, NY, USA.
- 8 Felsenstein Medical Research Center, Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel

Word count: 2,799

Corresponding Author:

Prof. Gal Shoval, MD
Geha Mental Health Center
P.O. Box 102, Petah Tiqva 49 100, Israel
Telephone: 972-3-925-8440, Fax: 972-3-925-8276; E-Mail: shovgal@tau.ac.il

Key words: antidepressants, mortality, adherence, anxiety, depression, Parkinson disease

Download English Version:

<https://daneshyari.com/en/article/8285768>

Download Persian Version:

<https://daneshyari.com/article/8285768>

[Daneshyari.com](https://daneshyari.com)