

www.elsevier.com/locate/redox

A reciprocal relationship between reactive oxygen species and mitochondrial dynamics in neurodegeneration

Clara Hiu-Ling Hung, Sally Shuk-Yee Cheng, Yuen-Ting Cheung, Suthicha Wuwongse, Natalie Qishan Zhang, Yuen-Shan Ho, Simon Ming-Yuen Lee, Raymond Chuen-Chung Chang

PII: S2213-2317(17)30503-7
DOI: http://dx.doi.org/10.1016/j.redox.2017.08.010
Reference: REDOX733

To appear in: *Redox Biology*

Received date: 10 July 2017
Revised date: 3 August 2017
Accepted date: 8 August 2017

Cite this article as: Clara Hiu-Ling Hung, Sally Shuk-Yee Cheng, Yuen-Ting Cheung, Suthicha Wuwongse, Natalie Qishan Zhang, Yuen-Shan Ho, Simon Ming-Yuen Lee and Raymond Chuen-Chung Chang, A reciprocal relationship between reactive oxygen species and mitochondrial dynamics in neurodegeneration, *Redox Biology,* http://dx.doi.org/10.1016/j.redox.2017.08.010

A reciprocal relationship between reactive oxygen species and mitochondrial dynamics in neurodegeneration

Clara Hiu-Ling Hung[1,3], Sally Shuk-Yee Cheng[1], Yuen-Ting Cheung[1], Suthicha Wuwongse[1], Natalie Qishan Zhang[1], Yuen-Shan Ho[4], Simon Ming-Yuen Lee[3], Raymond Chuen-Chung Chang[1,2*]

[1]Laboratory of Neurodegenerative Diseases, School of Biomedical Sciences, LKS Faculty of Medicine
[2]State Key Laboratory of Brain and Cognitive Sciences, The University of Hong Kong, Pokfulam, Hong Kong, China
[3]Institute of Chinese Medical Sciences, University of Macau, Macau, China
[4]School of Nursing, Faculty of Health and Social Sciences, The Hong Kong Polytechnic University, Hung Hom, Kowloon, Hong Kong, China

Correspondence to: Dr. Raymond Chuen-Chung CHANG, Laboratory of Neurodegenerative Diseases, School of Biomedical Sciences, LKS Faculty of Medicine, The University of Hong Kong. Rm. L4-49, Laboratory Block, Faculty of Medicine Building, 21 Sassoon Road, Pokfulam, Hong Kong SAR. Tel.: +852 3917 9127; Fax: +852 3917 0857. rccchang@hku.hk

Abstract

Mitochondrial fragmentation due to fission/fusion imbalance has often been linked to mitochondrial dysfunction and apoptosis in neurodegeneration. Conventionally, it is believed that once mitochondrial morphology shifts away from