


Accepted Manuscript

Loss of toll-like receptor 3 aggravates hepatic inflammation but ameliorates steatosis in mice

Young-Sun Lee, Doyeon Kim, Tae-Jun Kim, So Yeon Kim, Jae-Kwang Jung, Jae-Kap Choi, Hyon-Seung Yi, Jin-Seok Byun



PII: S0006-291X(18)30220-1

DOI: [10.1016/j.bbrc.2018.01.191](https://doi.org/10.1016/j.bbrc.2018.01.191)

Reference: YBBRC 39364


To appear in: *Biochemical and Biophysical Research Communications*

Received Date: 25 January 2018

Accepted Date: 31 January 2018

Please cite this article as: Y.-S. Lee, D. Kim, T.-J. Kim, S.Y. Kim, J.-K. Jung, J.-K. Choi, H.-S. Yi, J.-S. Byun, Loss of toll-like receptor 3 aggravates hepatic inflammation but ameliorates steatosis in mice, *Biochemical and Biophysical Research Communications* (2018), doi: 10.1016/j.bbrc.2018.01.191.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Graphical abstract. Suggested diagram of TLR3 mediated-changed TLR4 signaling is responsible for increased inflammation and decreased steatosis in TLR3 knockout mice.

Download English Version:

<https://daneshyari.com/en/article/8293754>

Download Persian Version:

<https://daneshyari.com/article/8293754>

[Daneshyari.com](https://daneshyari.com)