Accepted Manuscript



SIRT7 is an important regulator of cartilage homeostasis and osteoarthritis development

Wataru Korogi, Tatsuya Yoshizawa, Md Fazlul Karim, Hironori Tanoue, Masaki Yugami, Shihab U. Sobuz, Eiichi Hinoi, Yoshifumi Sato, Yuichi Oike, Hiroshi Mizuta, Kazuya Yamagata

PII: S0006-291X(18)30144-X

DOI: 10.1016/j.bbrc.2018.01.129

Reference: YBBRC 39302

To appear in: *Biochemical and Biophysical Research Communications*

Received Date: 13 January 2018

Accepted Date: 20 January 2018

Please cite this article as: W. Korogi, T. Yoshizawa, M.F. Karim, H. Tanoue, M. Yugami, S.U. Sobuz, E. Hinoi, Y. Sato, Y. Oike, H. Mizuta, K. Yamagata, SIRT7 is an important regulator of cartilage homeostasis and osteoarthritis development, *Biochemical and Biophysical Research Communications* (2018), doi: 10.1016/j.bbrc.2018.01.129.

SIRT7 is an important regulator of cartilage homeostasis and osteoarthritis development.

Wataru Korogi[1, 2], Tatsuya Yoshizawa[1], Md. Fazlul Karim[1], Hironori Tanoue[3], Masaki Yugami[3], Shihab U. Sobuz[1], Eiichi Hinoi[4], Yoshifumi Sato[1], Yuichi Oike[3], Hiroshi Mizuta[2], Kazuya Yamagata[1]

[1]*Department of Medical Biochemistry,* [2]*Department of Orthopaedic Surgery,* [3]*Department of Molecular Genetics, Faculty of Life Sciences, Kumamoto University, 1-1-1 Honjo, Chuo-ku, Kumamoto 860-8556, Japan.*

[4]*Laboratory of Molecular Pharmacology, Division of Pharmaceutical Sciences, Kanazawa University Graduate School of Natural Science and Technology, Kanazawa 920-1192, Japan.*

Running title: *SIRT7 is important for cartilage homeostasis.*

Correspondence to:

Tatsuya Yoshizawa, PhD

Department of Medical Biochemistry, Faculty of Life Sciences, Kumamoto University

1-1-1 Honjo, Kumamoto, Kumamoto 860-8556, Japan

yoshizaw@kumamoto-u.ac.jp

TEL: +81-96-373-5070

FAX: +81-96-364-6940