



Hyperosmolarity evokes histamine release from ileum mucosa by stimulating a cholinergic pathway

Banqin Wang^{a,1}, Ning An^{a,b,1}, Abdul Sami Shaikh^c, Haoyi Wang^a, Ling Xiao^a, Hongwei Liu^a, Jingxin Li^a, Dongbo Zhao^{b,*}

^a Department of Physiology, Shandong Provincial Key Laboratory of Mental Disorders, Shandong University School of Medicine, Jinan, China

^b Department of Thoracic Surgery, Shandong Cancer Hospital and Institute, Shandong Academy of Medical Sciences, Jinan, China

^c Institute of Clinical Pharmacology, Qilu Hospital of Shandong University, Jinan, China

ARTICLE INFO

Article history:

Received 11 September 2017

Accepted 17 September 2017

Available online 19 September 2017

Keywords:

Hyperosmolarity

Histamine

Ileum mucosa

Short-circuit current

Enteric nervous system

ABSTRACT

Changes in extracellular osmolarity lead to alteration in cellular volume. In the study, we examined the effects of hyperosmolarity on short-circuit currents (I_{SC}) in the rat ileum using the Ussing chamber technique. Mucosal exposure to 20 mM glucose evoked a decrease of I_{SC} in the rat ileum, which was antagonized by the stretch-activated channel blocker GdCl₃, TTX and atropine, respectively. In contrast, it was not blocked by phlorizin, a Na⁺-glucose cotransporter SGLT1 inhibitor. Furthermore, the unabsorbed substances, such as sucrose, lactulose or urea, also induced a decrease of I_{SC} in rat ileum. ELISA results revealed that 20 mM glucose stimulated the release of histamine from rat ileum mucosa, which was attenuated by TTX. In addition, the glucose-induced I_{SC} was depressed by pyrilamine, a histamine H₁ receptor blocker (H₁ antagonist) whereas it was not affected by ranitidine (H₂ antagonist), clobenpropit (H₃ antagonists) or JNJ777120 (H₄ antagonist), respectively. The ion substitution experiments suggest that the changes of Na⁺ and HCO₃⁻ ion flux underlie the glucose-induced I_{SC} . In conclusion, osmotic stimulus decreased the basal I_{SC} of rat ileum by evoking histamine release from ileum mucosa. The changes of Na⁺ and HCO₃⁻ ion transport are involved in the glucose-evoked decrease of basal I_{SC} .

© 2017 Elsevier Inc. All rights reserved.

1. Introduction

The intestinal epithelium is one of the first interfaces between the organism and the environment. During clinical manifestation, it delivers a selective permeable barrier that causes limitations to the permeability of luminal noxious molecules, for example toxins, pathogens and antigens, while permitting the necessary absorption of nutrients and water since maintenance of cell volume is crucial for this selective barrier. Disturbances in tonicity (effective osmolarity) are the major clinical disorders affecting cell volume [1]. Cell shrinking, secondary to hypertonicity, causes severe clinical manifestations and even death [2]. To date, there are several unresolved

aspects of cell volume regulation. These include the identity of the structure of the osmotic sensor, and the nature of signaling pathways of the hypothetical cell volume sensor which leads to activation of volume-dependent ion transporters. It has been reported that hypertonicity stimulates Cl⁻ transport in eel intestinal epithelium by the activation of the luminal Na⁺-K⁺-2Cl⁻ cotransporter (NKCC) and the functionality of basolateral Cl⁻ channels [3]. The present study reveals a new signaling pathway of cell volume regulation. We demonstrated that hyperosmolarity evoked histamine release from ileum mucosa, and then regulated Na⁺ and HCO₃⁻ transport to alter cell volume.

2. Materials and methods

2.1. Animals and tissue preparation

All prospective procedures throughout the experiment were followed by protocol and conducted with the procedures for the Care and Use of Laboratory Animals of Shandong University, and the work was approved by the Medical Ethics Committee for

Abbreviations: I_{SC} , short-circuit current; TTX, tetrodotoxin; NKCC, Na⁺-K⁺-2Cl⁻ cotransporter; ENS, enteric nervous system; ACh, acetylcholine; PLC, phospholipase C; IP3, inositol trisphosphate; DAG, diacylglycerol; NO, nitric oxide; AC, adenylyl cyclase.

* Corresponding author.

E-mail address: 1172296684@qq.com (D. Zhao).

¹ These authors contributed equally to this work.

Experimental Animals, Shandong University, China (number ECAESDUSM 2012029). The usage of adult male Wistar rats ranging between 200 and 250 g were employed in this study. This specie of rat was acquired from the Animal Center of Shandong University, China. Animals were allocated and allowed free access to water, had been fasted overnight prior before experiments and were then anesthetized and decapitated. Tissue preparations were done according to the previously described procedures [4]. After eliminating luminal contents smoothly, the segments of the ileum were chopped along the mesenteric border and tissues were pinned flat on a Sylgard-lined petri dish with the mucosal surface facing down. The serosa and muscularis were removed gently and precisely in order to get mucosal-submucosal preparations. Tissue preparations were unremittingly oxygenated with a gas mixture (95% O₂ and 5% CO₂) while being bathed in ice-cold Krebs solution during preparation. The composition of Krebs solution was (in mM): 120.6 NaCl, 5.9 KCl, 2.5 CaCl₂, 1.2 KH₂PO₄, 1.2 MgCl₂, 15.4 NaHCO₃ and 11.5 Glucose.

2.2. Short-circuit current (I_{SC}) measurement

I_{SC} in vitro was measured using Ussing Chamber techniques. The tissue preparations were attached and placed between 2 halves of the Ussing Chambers. The exposed area (about 0.50 cm²) was fortified with water-jacketed gas lifts which were then bathed on both sides with 5 mL Krebs solution and oxygenated with a gas mixture of 95% O₂ and 5% CO₂, while maintaining the pH at 7.4.

The Krebs solution was maintained at 37 °C and circulated through a reservoir. The tissue was then continuously voltage-clamped to a zero potential difference by using an external current with compensation for fluid resistance in consideration. The baseline value of the electrical parameters was measured as the mean over the 3 min interval immediately prior to administration of drug. The tissue was then equilibrated under these stated conditions for 20 min to maintain the I_{SC} proceeding to the addition of drugs. The transepithelial potential difference for each prospective preparation was calculated with Ag/AgCl reference electrodes (P2020S; Physiologic Instruments, San Diego, Calif) connected to a preamplifier that was in turn connected to a voltage clamp amplifier (VCC MC4; Physiologic Instruments, San Diego, Calif). The changes in the short circuit current (ΔI_{SC}) were calculated, and results were recorded on the basis of the value before and after the process of stimulation. The ΔI_{SC} was normalized as the current per unit area of epithelium ($\mu A/cm^2$). The viability of tissues was checked by inducing stimulation by acetylcholine (ACh).

2.3. Measurement of histamine

The ileum mucosa was attached between the 2 halves of the Ussing chambers (0.50 cm²). The tissues were there after bathed on both sides with 5 mL Krebs solution, gassed with 95% O₂ and 5% CO₂, pH adjusted to 7.4 and maintained persistently at 37 °C by circulating the solution of contents via a reservoir throughout the experiments. Glucose (20 mM) was added to the mucosa side solution then the tissues were taken out and homogenized and centrifuged for ELISA at 4 °C. Histamine was measured using a Histamine ELISA Kit for mouse (R&D, USA) with a detection range from 0.5 to 200.0 ng/mL. The histamine standard curve was calculated using a 4-PL curve fit ($R^2 = 0.9950$).

2.4. Drugs

The D-glucose (20 mM), phlorizin (100 μM), D (+)-Sucrose (20 mM), lactulose (20 mM) or saccharin (1 mM) was added to the mucosal side bathing solution and measured the changes of I_{SC} .

Atropine (10 μM) was used to block the cholinergic system. GdCl₃ (100 μM) was used to block the osmotic sensors. Histamine receptor antagonists including pyrilamine (H₁ antagonist, 1 μM), ranitidine (H₂ antagonist, 1 μM), clobenpropit (H₃ antagonist, 1 μM) or JNJ7777120 (H₄ antagonist, 1 μM) was added to serosal side bathing solution to study the effect of histamine to the hyperosmolarity. Tetrodotoxin (TTX, 1 μM) was used to block the neural pathway. Atropine, TTX or furosemide (100 μM) was added to the serosal side bathing solution. In the alternative experiments, the Cl⁻ free solution contained (in mM): 117 Na-gluconate, 4.7 K-gluconate, 8 Ca-(gluconate)₂, 1.2 Mg-(gluconate)₂, 1.2 NaH₂PO₄, 25 NaHCO₃, and 11 glucose. In HCO₃⁻-free solution, HCO₃⁻ was replaced by gluconate and HEPES buffer and the pH were adjusted to 8.0. In Na⁺-free solution, 120.6 mM NaCl was replaced by 120.6 mM KCl, pH 7.4 adjusted with KOH.

2.5. Data analysis and statistics

Data showed as means $\pm$ SEM and the n values represent the numbers of animals used in these experiments. We considered one-way ANOVA or unpaired Student's *t*-tests to investigate if there were significant differences in basal electrical parameters among the tissue elements. $P < 0.05$ was considered statistically significant.

3. Results

3.1. Osmotic stimulus evoked a decrease in I_{SC} in rat ileum

It's well known the absorption of glucose in intestinal epithelium causes an increase of I_{SC} because of Na⁺ co-transport. Our experiments confirmed that indeed, the addition of glucose (20 mM) to mucosal bathing solution suggested an increase in I_{SC} in rat jejunum (Fig. 1A). In contrast, 20 mM glucose evoked a decrease of I_{SC} in the rat ileum (Fig. 1B). The absorption of glucose in intestinal epithelium is through Na⁺ -glucose cotransporter SGLT1 therefore absorption of glucose is Na⁺ -dependent. Since phlorizin inhibits SGLT1, we added phlorizin to test whether the change in I_{SC} is evoked by SGLT1. Phlorizin blocked glucose-evoked increase of I_{SC} in jejunum, however, did not exert any effect on glucose-evoked decrease of I_{SC} in the rat ileum (Fig. 1C). Furthermore, sucrose or lactulose which is not absorbed by the small intestine also caused a decrease of I_{SC} (Fig. 1D and E). In addition, a non-nutrient osmotic load such as urea (20 mM) showed similar result (Fig. 1F). It is well known that the small intestine expresses sweet taste receptors [5,6]. To elucidate the role of sweet taste receptors, saccharin (1 mM), a sweet taste receptor agonist was used. However, it had no effect on the I_{SC} of the ileum mucosa (Fig. 1G). Notably, the stretch-activated channel blocker GdCl₃ (100 μM) largely attenuated the osmotic stimulus-evoked I_{SC} in the rat ileum (Fig. 1H). These results suggest that 20 mM glucose-induced I_{SC} in the rat ileum may be due to the hyperosmotic challenges.

3.2. The osmotic stimulus-evoked I_{SC} in the rat ileum is mediated by the cholinergic circuitry of enteric nervous system (ENS)

The ENS plays a significant part in the regulation of intestinal epithelial ion transport. The principal role of submucosal plexus is in sensing the environment within the lumen. To examine the contribution of the ENS in the effects of hypertonic challenge, TTX (10⁻⁶ M) was added to the serosal bathing solution to block the ENS. In the presence of TTX, the 20 mM glucose-induced I_{SC} was almost suppressed (Fig. 2A). Interestingly atropine, a selective muscarinic ACh receptor antagonist also inhibited the effect of 20 mM glucose on the I_{SC} of ileum mucosa (Fig. 2B).

Download English Version:

<https://daneshyari.com/en/article/8296296>

Download Persian Version:

<https://daneshyari.com/article/8296296>

[Daneshyari.com](https://daneshyari.com)