Accepted Manuscript



A soluble form of the interleukin-6 family signal transducer gp130 is dimerized via a C-terminal disulfide-bridge resulting from alternative mRNA splicing

Janina Wolf, Georg H. Waetzig, Torsten M. Reinheimer, Jürgen Scheller, Stefan Rose-John, Christoph Garbers

PII: S0006-291X(16)30127-9

DOI: 10.1016/j.bbrc.2016.01.127

Reference: YBBRC 35244

To appear in: *Biochemical and Biophysical Research Communications*

Received Date: 11 January 2016

Accepted Date: 21 January 2016

Please cite this article as: J. Wolf, G.H. Waetzig, T.M. Reinheimer, J. Scheller, S. Rose-John, C. Garbers, A soluble form of the interleukin-6 family signal transducer gp130 is dimerized via a C-terminal disulfide-bridge resulting from alternative mRNA splicing, *Biochemical and Biophysical Research Communications* (2016), doi: 10.1016/j.bbrc.2016.01.127.

A soluble form of the interleukin-6 family signal transducer gp130 is dimerized via a C-terminal disulfide-bridge resulting from alternative mRNA splicing

Janina Wolf[1], Georg H. Waetzig[2], Torsten M. Reinheimer[3], Jürgen Scheller[4], Stefan Rose-John[1], and Christoph Garbers[1,§]

[1]Institute of Biochemistry, Kiel University, Kiel, Germany;

[2]CONARIS Research Institute AG, Kiel, Germany;

[3]Non-Clinical Development, Ferring Pharmaceuticals A/S, Copenhagen, Denmark;

[4]Institute of Biochemistry and Molecular Biology II, Medical Faculty, Heinrich-Heine University, Düsseldorf, Germany

[§]Correspondence to: Dr. Christoph Garbers, Institute of Biochemistry, Kiel University, Rudolf-Hoeber-Strasse 1, 24118 Kiel, Germany; Tel: 0049-4318801676, Fax: 0049-4318805007; E-Mail: cgarbers@biochem.uni-kiel.de

Short title: Dimer formation of sgp130